



Annual Fisheries Statistics Report
for
Abu Dhabi Emirate 2007
(Project no. 02 – 23 – 0008 – 07)

Marine Environment Research Center
Environment Agency – Abu Dhabi

Contents

| | |
|--|----------|
| 1. Executive Summary | 3 |
| 2. Introduction | 4 |
| 3. Methodology and Survey Design | 5 |
| 4. Tables | |
| 4.1 Catch and Species Composition | |
| Table 1 - Total catch (tonnes) and species composition for the Emirate of Abu Dhabi..... | 9 |
| Table 2 - Total catch (tonnes) by landing site in the Emirate of Abu Dhabi..... | 12 |
| Table 3 - Total catch (tonnes) by boat/gear type in the Emirate of Abu Dhabi..... | 15 |
| Table 4 (a) - Total catch (tonnes) and species composition for landings at Sila..... | 18 |
| Table 4 (b) - Total catch (tonnes) and species composition for landings at Delma Island..... | 21 |
| Table 4 (c) - Total catch (tonnes) and species composition for landings at Al Marfa..... | 25 |
| Table 4 (d) - Total catch (tonnes) and species composition for landings at Al Bateen..... | 27 |
| Table 4 (e) - Total catch (tonnes) and species composition for landings at Free Port..... | 33 |
| Table 4 (f) - Total catch (tonnes) and species composition for landings at Al Sadar..... | 38 |
| 4.2 Effort | |
| Table 5 - Total number of trips by boat/gear type in the Emirate of Abu Dhabi..... | 41 |
| Table 6 - Total number of trips by landing site in the Emirate of Abu Dhabi..... | 42 |
| Table 7 (a) - Total number of boatdays for lansh by boat/gear type in the Emirate of Abu Dhabi..... | 43 |
| Table 7 (b) - Total number of boathours for tarad by boat/gear type in the Emirate of Abu Dhabi..... | 43 |
| Table 8 (a) - Total number of boatdays for lansh by landing site in the Emirate of Abu Dhabi..... | 44 |
| Table 8 (b) - Total number of boathours for tarad by landing site in the Emirate of Abu Dhabi..... | 45 |
| 4.3 Catch per Unit Effort (CPUE) | |
| Table 9 (a) - CPUE in kg/boatday for lansh landings at Sila..... | 46 |
| Table 9 (b) - CPUE in kg/boatday for lansh landings at Delma Island..... | 47 |
| Table 9 (c) - CPUE in kg/boatday for lansh landings at Free Port..... | 49 |
| Table 10 (a) - CPUE in kg/boathour for tarad landings at Sila..... | 52 |
| Table 10 (b) - CPUE in kg/boathour for tarad landings at Delma Island..... | 55 |
| Table 10 (c) - CPUE in kg/boathour for tarad landings at Marfa..... | 58 |
| Table 10 (d) - CPUE in kg/boathour for tarad landings at Al Bateen..... | 60 |
| Table 10 (e) - CPUE in kg/boathour for tarad landings at Free Port..... | 66 |
| Table 10 (f) - CPUE in kg/boathour for tarad landings at Al Sadar..... | 72 |

4.4 Wholesale value

| | |
|---|----|
| Table 11 - Wholesale value (AED '000) of landings by boat/gear type. | 75 |
| Table 12 - Wholesale value (AED '000) of landings by landing site. | 78 |

5. Figures

5.1 Catch and Species Composition

| | |
|--|----|
| Figure 1 – Total catch landed by landing site in the Emirate of Abu Dhabi. | 81 |
| Figure 2 - Total catch landed by month and boat type..... | 82 |
| Figure 3 - Total catch of principal species families landed by boat/gear type combination..... | 83 |

5.2 Effort

| | |
|--|----|
| Figure 4 (a) - Breakdown of leading gear type used on a Lansh in the Emirate of Abu Dhabi..... | 84 |
| Figure 4 (b) - Breakdown of leading gear type used on a Tarad in the Emirate of Abu Dhabi..... | 85 |
| Figure 5 - Total number of fishing trips by site and boat/gear type combination. | 86 |
| Figure 6 - Total number of fishing trips by month and boat/gear type combination..... | 87 |
| Figure 7 (a) - Mean duration of lansh/gargoor fishing trips (\pm 95% CI) by landing site..... | 88 |
| Figure 7 (b) - Mean duration of tarad/hadaq fishing trips (\pm 95% CI) by landing site | 89 |

5.3 Catch per Unit Effort (CPUE)

| | |
|---|----|
| Figure 8 (a) - CPUE (Kg/Trip) of principal species families from lansh/gargoor fisheries..... | 90 |
|---|----|

5.4 Wholesale value

| | |
|---|----|
| Figure 9 - Wholesale value of principal species families by boat/gear type combination..... | 93 |
| Figure 10 - Wholesale value of principal species families by landing site. | 94 |

6. Annexes

| | |
|---|----|
| Annex 1 - Landing sites and fishing grounds in the Emirate of Abu Dhabi..... | 95 |
| Annex 2 - Commercial species used in the catch and effort data recording system. | 96 |

1. Executive Summary

This report presents the fisheries statistics for Abu Dhabi during 2007 and provides a brief description of the fisheries of the Emirate and the survey design and methodology employed in their monitoring. The data presented contribute to EAD's strategic goal of developing a management regime for the fisheries of Abu Dhabi and are produced through the 'Fish Landings and Population Dynamics Project' of the Marine Environment Research Center.

An estimated 5,337 tonnes of fish was landed in the Emirate during 2007, 80 % of which consisted of demersal species. The total catch represented a decrease of 8 % from the previous year. Landings from lansh increased by 6.4 % to 3,029 tonnes, and landings from tarads dropped by 22.7 % to 2,308 tonnes compared to 2006. The most important species categories by weight were Groupers (23.7 %), Emperors (22.5 %), Jacks (16.8 %), Sweetlips (13.5 %) and Kingfish (8.1 %).

The majority of the catch (3,730 tonnes) was landed in Eastern Abu Dhabi, against 1,606 tonnes in Western Abu Dhabi. The Free Port remained the largest fish-landing site accounting 53.8 % of the total catch, followed by Delma Island (17.0 %), Marfa (9.0 %) and Al Bateen (8.8 %). Traditional wooden dhows or lansh deploying gargoor (fish traps) landed 3,029 tonnes, equivalent to 56.8 % of the total catch. Open fiberglass boats or tarads operating hand line (hadaq) landed 1,728 tonnes (32.4 %), followed by encircling nets (halaq) landing 205 tonnes (3.8 %) and encircling nets (Al Defara) landing 194 tonnes equivalent to 3.6 % of the total catch.

Total fishing effort during 2007 was estimated at 27,380 tarad trips and 6,266 lansh trips. The number of lansh trips increased by 2 % compared to 5,972 trips in 2006. In contrast, the number of tarad fishing trips decreased by 6 % compared to the previous year. The drop in tarad fishing trips was notable in all landing sites.

The total ex-vessel value of the landed catch increased by 4.1 % to 63.2 million AED, compared to 2006. By value, the most important species categories were Groupers (38.4 %), Kingfish (12.5 %), Jacks (14.3 %) and Emperors (19.4 %). Eastern Abu Dhabi accounted for 77 % of the value of the total landings, the most important site being the Free Port. By boat type, dhows accounted for 55.7 % of the value of the total landed catch.

2. Introduction

Fisheries provide a source of income, employment and recreation to inhabitants while contributing to the cultural heritage of the Emirate of Abu Dhabi. Fishing is primarily conducted from open fiberglass dories and wooden dhows, dome shaped traps are the most commonly used gear type although a variety of other methods exist. Catches are typically diverse and characteristic of multi-species tropical fisheries, target species primarily being composed of representatives of the families: Carangidae, Lethrinidae, Haemulidae, Serranidae and Scombridae.

Catch and effort data recording surveys form the foundation of fisheries management information systems. The outputs enable resource status to be monitored and evaluations of fisheries performance to be made, aspects crucial for informed management planning and decision-making. In this context, the Marine Environment Research Center of the Environment Agency – Abu Dhabi (EAD) is implementing the 'Fish Landings and Population Dynamics Project' the principal objectives of which include the collection of catch, effort and economic data and the production of fisheries statistics for the Emirate of Abu Dhabi. The project responds directly to the EAD's goal of developing a management regime for the fisheries of Abu Dhabi.

Since the inception of the Fish Landings and Population Dynamics Project in 2001, a stratified catch and effort data recording system was set in place. Starting 2005, an improved fishery data collection system was launched with a higher resolution in the recordings of species landed and gear types used. In addition to valuable input by key stakeholder organizations such as Coast Guard, Abu Dhabi Fishermen Cooperative Society and Delma Cooperative Society, was the level of cooperation between the Ministry of Environment and Water and EAD very instrumental regarding the collection of data by enumerators. Alongside this upgrade, a new tailor-made fisheries database system was developed capable of producing instantaneous fisheries statistical reports.

This report presents the fisheries statistics for Abu Dhabi for 2007 and provides a brief description of the fisheries of the Emirate and the survey methodology employed in their monitoring.

3. Methodology and Survey Design

Landing site stratification: For estimation purposes, the Emirate of Abu Dhabi is divided into two strata namely: Eastern Abu Dhabi and Western Abu Dhabi. There are eight fish landing sites in the Emirate: Sila, Delma Island, Jebel Dhanna, Marfa, Al Bateen, Free Port, Al Sadar, and Ghantoot (see Annex 1).

Jebel Dhanna is an inactive landing site since mid 2004. As the physical infrastructure still enables Jebel Dhanna to re-commence as a commercial landing site at any time, it is not excluded from the fisheries statistics presented.

Ghantoot (formerly known as Khor Al Azab) is a landing site used by recreational fishermen and thus not covered by present data collection system of commercial fisheries. However, the entry Ghantoot still appears in statistical tables below only to maintain continuity with the 2004 statistical reports where Ghantoot landings were, erroneously presented as that of commercial fisheries.

Fishing boats stratification: There are two principal fishing boat types in the Emirate of Abu Dhabi, the tarad and lansh. The tarad is an open dory usually of fiberglass construction powered by an outboard motor/s, fishing trips are conducted during the day or the night depending on the gear type being used and target species. Due to the small size of these boats, trip duration usually does not exceed 24 hours. There are about 773 tarads licensed in Abu Dhabi (January 2008). The lansh is a decked boat usually constructed of wood and powered by an in-board engine/s, the average trip duration usually ranging between 1 and 7 days. There are about 339 Lansh licensed in Abu Dhabi (January 2008).

Fishing gear stratification: The fisheries in the Emirate of Abu Dhabi are multi-gear, given the many different fishing gears that are often used simultaneously during fishing trips.

Due to the indirect manner in how effort data is collected, it is not possible to discern all possible gear type combinations and sort out the proportionate landing compositions. As such, the statistical tables presented refer to the leading gear type in a combination of gears being used. For instance, in the case of a lansh deploying gargoor plus hadaq, the leading gear type is gargoor, as shown in the overview below.

| <u>Leading gear</u> | <u>Operated by</u> | <u>Description</u> |
|---------------------|--------------------|--|
| <u>Gargoor</u> | Lansh only | Dome shaped wire traps ($\geq 1.5 \text{ m}^3$), usually baited with dried sardines and/or bread and used to target demersal species, such as Hamour, Shaari, Jesh and Safi. <i>Used alone or in combination with Hadaq and/or Leikh.</i> |
| <u>Leikh</u> | Lansh or Tarad | Gill nets up to 200 m in length and 6m deep set at the surface and used to target large pelagic species, such as Kanaad, Dhill'e and Jesh. <i>Used alone or in combination with Hadaq.</i> |
| <u>Hadaq</u> | Lansh or Tarad | Monofilament hand line with one or more baited hooks, targeting demersal species such as Hamour and Shaari. <i>Used alone.</i> |
| <u>Al Defara</u> | Tarad only | Encircling gillnets up to 500 meters long and 4 meters high set in shallow coastal waters targeting demersal species such as Jesh and Safi. <i>Used alone or in combination with Hadaq, Leikh and/or Halaq.</i> |
| <u>Leikh</u> | Tarad only | Small monofilament gillnets of 20 meters wide set in shallow water near the shore and used to target small demersal species, such as Naiser and Qabqoob <i>Used alone or in combination with Hadaq.</i> |
| <u>Halaq</u> | Tarad only | Encircling gillnets set around schools of pelagic species, such as Kanaad. <i>Used alone or in combination with Hadaq</i> |
| <u>Al Sakkar</u> | Tarad only | Barrier nets up to 1500 meters long and 4 meters high set at high tide and used to shut off a portion of the seabed, trapping small demersal species such as Qabit and Bedah behind the net as the tide drops. <i>Used alone.</i> |
| <u>Al Hadhra</u> | Tarad only | Fixed coral shaped enclosure trap set in the intertidal zone and connected to the shore via a long fence set perpendicular to the coastline. During ebb tide, the fence will guide fish into the entrance of the enclosure. Main target species is Naiser and Aifah. <i>Used alone.</i> |

Species classification: The gears operated in the different fisheries of the Emirate of Abu Dhabi catch over 100 species from more than 35 families. Landing data are recorded to the species level or, in some cases, to an unidentified family member (spp.). Annex 1 gives a detailed list of the main commercial species found in the fisheries of the Emirate of Abu Dhabi.

Data sources: Catch data was obtained from fish landing receipts issued by the Abu Dhabi/Delma Fishermen's Cooperative Society for the purchase of entire landed catch. The minimum sample size was set to all landing receipts of which the landing date equals one of ten predetermined days per month per landing site. In those landing sites where no Fishermen's Cooperative Society was present or fish landing receipts were unsuitable, enumerators recorded catch data. A minimum sample target size for complete catch and effort records was set to 10 % of the total landings per month per landing site.

Fishing effort data was derived from the boat movement logbook records kept by the UAE Coast Guard in all landing sites, except Marfa and Al Bateen, during 2007. Minimum sample size was set to all boat movement records of which the departure date equals one of ten predetermined days per month per landing site. At those landing sites for which no Coast Guard boat movement logbook records are available, the effort was estimated by tallying moored boat counts in Marfa, Al Bateen and in Sila, as explained further in Hartmann, et al (2007)¹.

The fish price per species was obtained from the Abu Dhabi Fishermen's Cooperative Society receipts for 2007.

Explanation of estimations: Total effort is estimated in number of boatdays for a lanch and number of boathours for a tarad fishing trip by landing site, boat and gear type combination on a monthly basis for 2007.

The Catch per Unit of Effort (CPUE) is estimated as kg/boatdays for lanch or as kg/boathours for tarad landings by landing site, boat and gear type combination and by family and species on a monthly basis for 2007.

Total catch is estimated by raising the CPUE with the total effort by landing site, boat and gear type combination, and family and species on a monthly basis for 2007, as described further in Hartmann, et al (2007)¹.

¹ Hartmann S., Al Abdessalaam T., Grandcourt E., Al Shamsi A., Francis F., Al Ali K., Al Ali S., Al Suwaidi A., 2007. **Design of INZAL ASMAK Emirates Fisheries Database System.** MERC, EAD. Abu Dhabi (*unpub*). 29 p.

Explanation of tables and figures: CPUE comparisons between lanch gargoor landings in Delma Island and Free Port, representing Western and Eastern Abu Dhabi strata respectively, are expressed in kg per trip rather than kg per boatday. The reason behind this is twofold: trip duration (including travel time) between homeport and fishing grounds for Delma Island fishermen is only a fraction of that spent by Free Port fishermen, and gargoor being a passive gear; the catch quantity is not related to trip duration.

Tables and figures are self-explanatory though it should be noted that any discrepancy between totals and the sum of components is due to rounding.

Table 1 - Total catch (tonnes) and species composition for the Emirate of Abu Dhabi.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|---------------------------|-------------------|-------------|-------------|-------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|--------------|-------------|
| Ariidae | | | 2.7 | 4.5 | 0.2 | 6.0 | 2.6 | 4.2 | 0.5 | 0.6 | 0.0 | 0.3 | 1.7 | 5.6 | 28.9 | 0.5 |
| | Arius thalassinus | Khan | 2.7 | 4.5 | 0.2 | 6.0 | 2.6 | 4.2 | 0.5 | 0.6 | 0.0 | 0.3 | 1.7 | 5.6 | 28.9 | 0.5 |
| Belonidae | | | 0.3 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| | Belonidae et al. | Haqool | 0.3 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| Carangidae | | | 97.3 | 84.1 | 97.6 | 77.7 | 110.9 | 86.2 | 55.8 | 41.2 | 56.2 | 65.8 | 73.5 | 50.8 | 896.9 | 16.8 |
| | Alepes vari | Gamlouh | 0.1 | 0.0 | 0.0 | 0.1 | 0.2 | 0.3 | 0.4 | 0.2 | 0.0 | 0.3 | 0.1 | 0.1 | 1.8 | 0.0 |
| | Atule mate | Durduman | 15.5 | 13.5 | 17.1 | 10.8 | 38.0 | 25.2 | 5.5 | 9.8 | 11.1 | 21.5 | 12.0 | 5.7 | 185.7 | 3.5 |
| | Carangidae all | Jesh unidentified | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Carangoides bajad | Jesh Um Al Hala | 13.9 | 17.1 | 18.6 | 42.1 | 14.9 | 19.1 | 21.5 | 10.3 | 10.3 | 14.5 | 16.8 | 8.7 | 207.8 | 3.9 |
| | Carangoides fulvoguttatus | Jesh Zeria | 40.3 | 37.8 | 24.0 | 2.7 | 16.8 | 3.3 | 5.0 | 1.2 | 0.3 | 10.6 | 20.3 | 16.3 | 178.5 | 3.3 |
| | Carangoides malabaricus | Jesh Sal | 2.0 | 0.5 | 0.0 | 4.6 | 2.4 | 2.1 | 0.1 | 0.2 | 11.2 | 0.0 | 6.3 | 1.0 | 30.5 | 0.6 |
| | Caranx heberi | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.5 | 0.9 | 1.0 | 3.1 | 1.5 | 0.3 | 0.6 | 9.1 | 0.2 |
| | Caranx ignobilis | Gufdar | 1.9 | 0.5 | 1.1 | 0.9 | 4.7 | 9.1 | 2.4 | 5.1 | 1.3 | 1.4 | 2.4 | 2.1 | 33.0 | 0.6 |
| | Gnathanodon speciosus | Zuraidi | 21.1 | 5.0 | 18.2 | 11.0 | 22.4 | 17.3 | 11.3 | 8.7 | 10.6 | 10.3 | 7.3 | 12.9 | 156.0 | 2.9 |
| | Megalaspis cordyla | Diyayuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.0 |
| | Parastromateus niger | Halwayoh | 0.0 | 0.1 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.8 | 0.0 |
| | Scomberoides | Dhil'e | 0.1 | 7.5 | 5.2 | 1.4 | 5.1 | 2.0 | 1.2 | 0.8 | 2.5 | 5.0 | 0.9 | 0.4 | 32.2 | 0.6 |
| | Scomberoides tol | Bassar | 0.5 | 0.5 | 12.1 | 2.6 | 1.0 | 1.8 | 6.5 | 2.6 | 2.0 | 0.7 | 6.7 | 1.1 | 38.2 | 0.7 |
| | Seriola dumerili | Jib | 0.0 | 1.2 | 0.6 | 1.3 | 5.0 | 4.4 | 1.0 | 1.2 | 3.7 | 0.1 | 0.0 | 0.6 | 19.1 | 0.4 |
| | Seriolina nigrofasciata | Hamam | 1.6 | 0.4 | 0.3 | 0.1 | 0.2 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.8 | 3.6 | 0.1 |
| Carcharhinidae | | | 1.1 | 1.8 | 0.2 | 0.4 | 0.4 | 0.0 | 0.1 | 0.4 | 0.0 | 8.7 | 0.0 | 0.1 | 13.2 | 0.2 |
| | Carcharhinidae et al. | Jarjoor | 1.1 | 1.8 | 0.2 | 0.4 | 0.4 | 0.0 | 0.1 | 0.4 | 0.0 | 8.7 | 0.0 | 0.1 | 13.2 | 0.2 |
| Chanidae | | | 1.8 | 0.1 | 0.1 | 0.0 | 2.7 | 0.0 | 1.3 | 4.6 | 0.0 | 0.0 | 0.0 | 0.0 | 10.6 | 0.2 |
| | Chanos chanos | Aifah | 1.8 | 0.1 | 0.1 | 0.0 | 2.7 | 0.0 | 1.3 | 4.6 | 0.0 | 0.0 | 0.0 | 0.0 | 10.6 | 0.2 |
| Clupeidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Sardinella longiceps | Oumah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ephippidae | | | 0.2 | 0.1 | 0.1 | 0.0 | 0.1 | 1.5 | 4.7 | 3.6 | 6.5 | 4.4 | 0.6 | 0.7 | 22.5 | 0.4 |
| | Ephippidae et al. | Imad | 0.2 | 0.1 | 0.1 | 0.0 | 0.1 | 1.5 | 4.7 | 3.6 | 6.5 | 4.4 | 0.6 | 0.7 | 22.5 | 0.4 |
| Gerreidae | | | 12.2 | 6.2 | 13.2 | 0.2 | 0.0 | 0.6 | 1.1 | 3.9 | 3.6 | 0.0 | 6.0 | 11.2 | 58.2 | 1.1 |
| | Gerres longirostris | Badah | 12.2 | 6.2 | 13.2 | 0.2 | 0.0 | 0.6 | 1.1 | 3.9 | 3.6 | 0.0 | 6.0 | 11.2 | 58.2 | 1.1 |
| Haemulidae | | | 41.4 | 27.8 | 58.6 | 127.3 | 63.4 | 69.7 | 55.5 | 50.4 | 41.2 | 51.4 | 61.1 | 71.8 | 719.7 | 13.5 |
| | Diagramma pictum | Farsh | 26.3 | 17.0 | 42.4 | 120.3 | 59.7 | 68.2 | 51.3 | 48.5 | 39.0 | 48.2 | 36.8 | 65.6 | 623.4 | 11.7 |
| | Plectorhinchus gaterinus | Hilali | 6.1 | 1.5 | 3.4 | 0.9 | 1.8 | 1.1 | 2.2 | 1.4 | 0.9 | 2.0 | 1.4 | 3.4 | 26.1 | 0.5 |
| | Plectorhinchus sordidus | Yanam | 9.0 | 9.3 | 12.7 | 4.0 | 1.9 | 0.4 | 0.7 | 0.4 | 0.5 | 1.2 | 0.7 | 2.4 | 43.2 | 0.8 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.2 | 2.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.8 | 0.0 | 22.2 | 0.5 | 27.0 | 0.5 |

Table 1 - Total catch (tonnes) and species composition for the Emirate of Abu Dhabi (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|--------------------------|-----------------------|-----------------|--------------|-------------|--------------|--------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|--------------|----------------|-------------|
| Labridae | | | 0.0 | 2.3 | 0.0 | 0.6 | 1.2 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 4.3 | 0.1 |
| | Choerodon robustus | Jin | 0.0 | 2.3 | 0.0 | 0.6 | 1.2 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 4.3 | 0.1 |
| Lethrinidae | | | 116.3 | 90.1 | 126.9 | 228.8 | 127.1 | 107.7 | 84.5 | 62.8 | 31.9 | 49.7 | 65.4 | 109.2 | 1,200.3 | 22.5 |
| | Lethrinus borbonicus | Yemah | 6.3 | 6.0 | 6.3 | 16.0 | 3.9 | 5.9 | 4.3 | 3.5 | 2.3 | 2.7 | 2.2 | 2.5 | 61.8 | 1.2 |
| | Lethrinus lentjan | Shaari Eshkheli | 4.5 | 3.9 | 8.6 | 21.0 | 12.0 | 10.9 | 15.2 | 14.1 | 3.7 | 11.5 | 6.9 | 8.8 | 121.0 | 2.3 |
| | Lethrinus microdon | Souli | 1.8 | 2.1 | 1.6 | 4.0 | 4.2 | 3.8 | 4.2 | 3.2 | 1.6 | 3.2 | 1.7 | 4.2 | 35.7 | 0.7 |
| | Lethrinus nebulosus | Shaari | 103.7 | 78.1 | 110.4 | 187.8 | 106.9 | 87.1 | 60.8 | 42.1 | 24.3 | 32.4 | 54.5 | 93.8 | 981.7 | 18.4 |
| Lutjanidae | | | 0.3 | 0.2 | 1.0 | 1.4 | 11.6 | 11.5 | 9.3 | 16.8 | 8.0 | 0.9 | 1.3 | 0.7 | 63.1 | 1.2 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.0 | 0.0 | 1.0 | 10.4 | 11.1 | 8.8 | 16.2 | 2.1 | 0.1 | 0.3 | 0.2 | 50.1 | 0.9 |
| | Lutjanus fulviflamma | Aqalah | 0.2 | 0.1 | 1.0 | 0.3 | 1.0 | 0.3 | 0.3 | 0.6 | 0.6 | 0.8 | 0.1 | 0.0 | 5.3 | 0.1 |
| | Lutjanus malabaricus | Hamra | 0.1 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 0.1 | 0.1 | 0.1 | 0.8 | 0.5 | 2.1 | 0.0 |
| | Lutjanus russelli | Umm Dhrais | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.4 | 0.0 |
| | Pinjalo pinjalo | Neimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.1 | 0.0 | 0.0 | 0.0 | 5.1 | 0.1 |
| Mixed | | | 0.6 | 0.5 | 0.6 | 1.4 | 1.3 | 0.6 | 1.6 | 1.0 | 0.4 | 0.4 | 0.2 | 1.3 | 9.8 | 0.2 |
| | Mixed species | Mekhlot | 0.6 | 0.5 | 0.6 | 1.4 | 1.3 | 0.6 | 1.6 | 1.0 | 0.4 | 0.4 | 0.2 | 1.3 | 9.8 | 0.2 |
| Mugilidae | | | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 3.5 | 0.2 | 1.2 | 0.0 | 0.0 | 59.8 | 3.6 | 70.1 | 1.3 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 3.5 | 0.2 | 1.2 | 0.0 | 0.0 | 59.8 | 3.6 | 70.1 | 1.3 |
| Myliobatidae etc. | | | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 |
| | Myliobatidea et al. | Lugmah | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 |
| Nemipteridae | | | 3.3 | 4.5 | 6.1 | 4.4 | 0.9 | 1.8 | 0.6 | 0.7 | 1.4 | 1.0 | 0.8 | 3.1 | 28.6 | 0.5 |
| | Scolopsis taeniatus | Ebzimi | 3.3 | 4.5 | 6.1 | 4.4 | 0.9 | 1.8 | 0.6 | 0.7 | 1.4 | 1.0 | 0.8 | 3.1 | 28.6 | 0.5 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.4 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.4 | 0.0 |
| Pomacanthidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Pomacanthus maculosus | Anfooz | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Portunidae | | | 12.5 | 23.8 | 19.6 | 37.7 | 1.7 | 1.5 | 1.1 | 3.2 | 1.8 | 8.8 | 2.8 | 8.7 | 123.3 | 2.3 |
| | Portunus pelagicus | Qabqoob | 12.5 | 23.8 | 19.6 | 37.7 | 1.7 | 1.5 | 1.1 | 3.2 | 1.8 | 8.8 | 2.8 | 8.7 | 123.3 | 2.3 |
| Rachycentridae | | | 2.5 | 3.2 | 3.0 | 2.3 | 5.0 | 1.8 | 0.6 | 1.3 | 0.1 | 2.7 | 1.5 | 2.7 | 26.8 | 0.5 |
| | Rachycentron canadum | Sikkil | 2.5 | 3.2 | 3.0 | 2.3 | 5.0 | 1.8 | 0.6 | 1.3 | 0.1 | 2.7 | 1.5 | 2.7 | 26.8 | 0.5 |
| Scaridae | | | 0.0 | 0.1 | 0.1 | 1.5 | 0.2 | 1.6 | 0.6 | 0.2 | 0.5 | 1.3 | 0.1 | 0.6 | 6.8 | 0.1 |
| | Scarus ghobban | Gain | 0.0 | 0.1 | 0.1 | 1.5 | 0.2 | 1.6 | 0.6 | 0.2 | 0.5 | 1.3 | 0.1 | 0.6 | 6.8 | 0.1 |

Table 1 - Total catch (tonnes) and species composition for the Emirate of Abu Dhabi (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|---------------------------|----------------|--------------|--------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|----------------|-------------|
| Scombridae | | | 84.0 | 134.3 | 74.7 | 1.2 | 10.6 | 5.0 | 6.3 | 1.4 | 57.8 | 36.7 | 17.6 | 4.1 | 433.7 | 8.1 |
| | Euthynnus affinis | Sadah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.7 | 0.0 |
| | Scomberomorus commerson | Kanaad | 84.0 | 134.3 | 74.7 | 1.2 | 10.1 | 4.9 | 6.3 | 1.4 | 57.8 | 36.7 | 17.5 | 4.1 | 432.9 | 8.1 |
| Scyllaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Thenus orientalis | Umm Ar Rubiyan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Sepiidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Sepia pharaonis | Habbar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Serranidae | | | 223.3 | 121.4 | 154.3 | 186.6 | 70.3 | 81.3 | 68.2 | 62.4 | 49.0 | 93.6 | 66.9 | 87.5 | 1,264.6 | 23.7 |
| | Cephalopholis hemistiktos | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Epinephelus areolatus | Semman | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Epinephelus coioides | Hamour | 223.3 | 121.4 | 154.3 | 186.5 | 70.2 | 81.1 | 68.1 | 62.4 | 49.0 | 93.5 | 66.9 | 87.5 | 1,264.1 | 23.7 |
| Siganidae | | | 0.4 | 10.3 | 0.9 | 4.3 | 1.9 | 1.0 | 2.7 | 2.0 | 1.5 | 2.0 | 0.6 | 3.6 | 31.3 | 0.6 |
| | Siganus canaliculatus | Safi Arabi | 0.4 | 10.3 | 0.9 | 4.3 | 1.9 | 1.0 | 2.7 | 2.0 | 1.5 | 2.0 | 0.6 | 3.6 | 31.3 | 0.6 |
| | Siganus javus | Safi Sneifi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sparidae | | | 48.3 | 20.8 | 22.6 | 13.7 | 8.3 | 4.3 | 3.4 | 9.9 | 23.9 | 61.8 | 6.6 | 14.9 | 238.4 | 4.5 |
| | Acanthopagrus bifasciatus | Faskar | 13.9 | 7.1 | 11.5 | 4.6 | 3.6 | 3.3 | 2.6 | 2.8 | 2.1 | 2.1 | 1.3 | 4.1 | 59.1 | 1.1 |
| | Acanthopagrus latus | Shaam | 0.5 | 0.5 | 1.4 | 0.8 | 0.2 | 0.4 | 0.7 | 0.1 | 0.0 | 0.3 | 0.0 | 0.3 | 5.4 | 0.1 |
| | Argyrops spinifer | Kofar | 14.3 | 7.6 | 9.4 | 8.0 | 4.4 | 0.6 | 0.1 | 0.0 | 0.1 | 0.0 | 0.2 | 1.8 | 46.4 | 0.9 |
| | Rhabdosargus sarba | Qabit | 19.6 | 5.7 | 0.2 | 0.2 | 0.0 | 0.0 | 0.0 | 7.0 | 21.7 | 59.4 | 5.0 | 8.8 | 127.5 | 2.4 |
| Sphyraenidae | | | 14.5 | 18.7 | 12.1 | 2.9 | 2.6 | 2.2 | 2.3 | 0.3 | 1.3 | 6.5 | 7.7 | 10.9 | 82.0 | 1.5 |
| | Sphyraena jello | Jedd | 14.5 | 18.7 | 12.1 | 2.9 | 2.6 | 2.2 | 2.3 | 0.3 | 1.3 | 6.5 | 7.7 | 10.9 | 82.0 | 1.5 |
| Stromateidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Pampus argenteus | Zubaidi | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.4 | 0.0 |
| Teraponidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Terapon jarbua | Yelly | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tetraodontidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Arothron stellatus | Faqel | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Total | | | 663.7 | 554.7 | 593.9 | 698.8 | 422.9 | 386.6 | 300.5 | 268.3 | 285.2 | 396.2 | 374.2 | 391.6 | 5,336.6 | 100.0 |
| (%) | | | 12.4 | 10.4 | 11.1 | 13.1 | 7.9 | 7.2 | 5.6 | 5.0 | 5.3 | 7.4 | 7.0 | 7.3 | 100.0 | |

Table 2 - Total catch (tonnes) by landing site in the Emirate of Abu Dhabi.

| Family | Species | Arabic name | Sila | Delma | Jebel Dhanna | Marfa | Al Bateen | Free Port | Al Sadar | Ghantoot | Total | (%) |
|-----------------------|---------------------------|-------------------|-------------|--------------|--------------|--------------|-------------|--------------|-------------|------------|--------------|-------------|
| Ariidae | | | 0.0 | 0.9 | 0.0 | 6.5 | 0.0 | 18.1 | 3.5 | 0.0 | 28.9 | 0.5 |
| | Arius thalassinus | Khan | 0.0 | 0.9 | 0.0 | 6.5 | 0.0 | 18.1 | 3.5 | 0.0 | 28.9 | 0.5 |
| Belonidae | | | 0.0 | 0.1 | 0.0 | 0.2 | 0.1 | 0.2 | 0.3 | 0.0 | 0.9 | 0.0 |
| | Belonidae et al. | Haqool | 0.0 | 0.1 | 0.0 | 0.2 | 0.1 | 0.2 | 0.3 | 0.0 | 0.9 | 0.0 |
| Carangidae | | | 34.3 | 91.9 | 0.0 | 150.4 | 28.8 | 518.5 | 73.0 | 0.0 | 896.9 | 16.8 |
| | Alepes vari | Gamlouh | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 1.6 | 0.0 | 0.0 | 1.8 | 0.0 |
| | Atule mate | Durduman | 0.2 | 17.1 | 0.0 | 16.3 | 0.6 | 145.0 | 6.6 | 0.0 | 185.7 | 3.5 |
| | Carangidae all | Jesh unidentified | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Carangoides bajad | Jesh Um Al Hala | 26.3 | 20.5 | 0.0 | 31.2 | 8.9 | 98.0 | 22.9 | 0.0 | 207.8 | 3.9 |
| | Carangoides fulvoguttatus | Jesh Zeria | 0.0 | 0.1 | 0.0 | 98.3 | 3.5 | 41.3 | 35.3 | 0.0 | 178.5 | 3.3 |
| | Carangoides malabaricus | Jesh Sal | 1.1 | 0.1 | 0.0 | 0.0 | 0.0 | 29.3 | 0.0 | 0.0 | 30.5 | 0.6 |
| | Caranx heberi | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.1 | 0.0 | 0.0 | 9.1 | 0.2 |
| | Caranx ignobilis | Gufdar | 1.0 | 14.1 | 0.0 | 0.0 | 0.0 | 17.9 | 0.0 | 0.0 | 33.0 | 0.6 |
| | Gnathanodon speciosus | Zuraidi | 0.5 | 31.1 | 0.0 | 0.1 | 0.0 | 124.3 | 0.1 | 0.0 | 156.0 | 2.9 |
| | Megalaspis cordyla | Diyayuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.3 | 0.0 |
| | Parastromateus niger | Halwayoh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.8 | 0.0 |
| | Scomberoides | Dhil'e | 0.0 | 6.5 | 0.0 | 0.4 | 8.6 | 10.9 | 5.8 | 0.0 | 32.2 | 0.6 |
| | Scomberoides tol | Bassar | 5.0 | 0.0 | 0.0 | 3.9 | 7.3 | 19.8 | 2.3 | 0.0 | 38.2 | 0.7 |
| | Seriola dumerili | Jib | 0.0 | 2.1 | 0.0 | 0.3 | 0.0 | 16.7 | 0.1 | 0.0 | 19.1 | 0.4 |
| | Seriolina nigrofasciata | Hamam | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 3.5 | 0.0 | 0.0 | 3.6 | 0.1 |
| Carcharhinidae | | | 1.9 | 0.8 | 0.0 | 0.0 | 7.7 | 2.5 | 0.3 | 0.0 | 13.2 | 0.2 |
| | Carcharhinidae et al. | Jarjoor | 1.9 | 0.8 | 0.0 | 0.0 | 7.7 | 2.5 | 0.3 | 0.0 | 13.2 | 0.2 |
| Chanidae | | | 1.9 | 0.0 | 0.0 | 0.0 | 8.6 | 0.1 | 0.0 | 0.0 | 10.6 | 0.2 |
| | Chanos chanos | Aifah | 1.9 | 0.0 | 0.0 | 0.0 | 8.6 | 0.1 | 0.0 | 0.0 | 10.6 | 0.2 |
| Clupeidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Sardinella longiceps | Oumah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ephippidae | | | 0.0 | 5.2 | 0.0 | 0.0 | 0.0 | 17.3 | 0.0 | 0.0 | 22.5 | 0.4 |
| | Ephippidae et al. | Imad | 0.0 | 5.2 | 0.0 | 0.0 | 0.0 | 17.3 | 0.0 | 0.0 | 22.5 | 0.4 |
| Gerreidae | | | 0.1 | 0.0 | 0.0 | 0.0 | 57.8 | 0.2 | 0.0 | 0.0 | 58.2 | 1.1 |
| | Gerres longirostris | Badah | 0.1 | 0.0 | 0.0 | 0.0 | 57.8 | 0.2 | 0.0 | 0.0 | 58.2 | 1.1 |
| Haemulidae | | | 60.0 | 304.8 | 0.0 | 5.8 | 24.7 | 319.7 | 4.7 | 0.0 | 719.7 | 13.5 |
| | Diagramma pictum | Farsh | 59.7 | 304.6 | 0.0 | 3.9 | 0.0 | 251.9 | 3.3 | 0.0 | 623.4 | 11.7 |
| | Plectorhinchus gaterinus | Hilali | 0.0 | 0.1 | 0.0 | 1.0 | 0.0 | 24.7 | 0.4 | 0.0 | 26.1 | 0.5 |
| | Plectorhinchus sordidus | Yanam | 0.3 | 0.2 | 0.0 | 0.0 | 0.4 | 42.3 | 0.0 | 0.0 | 43.2 | 0.8 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.9 | 24.3 | 0.9 | 0.9 | 0.0 | 27.0 | 0.5 |

Table 2 - Total catch (tonnes) by landing site in the Emirate of Abu Dhabi (cont.).

| Family | Species | Arabic name | Sila | Delma | Jebel Dhanna | Marfa | Al Bateen | Free Port | Al Sadar | Ghantoot | Total | (%) |
|--------------------------|-----------------------|-----------------|-------------|--------------|--------------|--------------|--------------|--------------|-------------|------------|----------------|-------------|
| Labridae | | | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.6 | 2.9 | 0.0 | 4.3 | 0.1 |
| | Choerodon robustus | Jin | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.6 | 2.9 | 0.0 | 4.3 | 0.1 |
| Lethrinidae | | | 11.3 | 204.0 | 0.0 | 117.4 | 7.4 | 779.2 | 81.0 | 0.0 | 1,200.3 | 22.5 |
| | Lethrinus borbonicus | Yemah | 0.4 | 2.2 | 0.0 | 6.4 | 0.1 | 50.1 | 2.6 | 0.0 | 61.8 | 1.2 |
| | Lethrinus lentjan | Shaari Eshkheli | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 119.4 | 0.9 | 0.0 | 121.0 | 2.3 |
| | Lethrinus microdon | Souli | 0.3 | 14.1 | 0.0 | 0.6 | 0.0 | 19.9 | 0.8 | 0.0 | 35.7 | 0.7 |
| | Lethrinus nebulosus | Shaari | 10.6 | 187.7 | 0.0 | 109.6 | 7.3 | 589.8 | 76.6 | 0.0 | 981.7 | 18.4 |
| Lutjanidae | | | 0.0 | 0.2 | 0.0 | 0.2 | 38.8 | 23.7 | 0.2 | 0.0 | 63.1 | 1.2 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.2 | 0.0 | 0.2 | 38.8 | 10.7 | 0.2 | 0.0 | 50.1 | 0.9 |
| | Lutjanus fulviflamma | Aqalah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.3 | 0.0 | 0.0 | 5.3 | 0.1 |
| | Lutjanus malabaricus | Hamra | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 0.0 | 0.0 | 2.1 | 0.0 |
| | Lutjanus russelli | Umm Dhrais | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Pinjalo pinjalo | Neimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.1 | 0.0 | 0.0 | 5.1 | 0.1 |
| Mixed | | | 0.3 | 0.0 | 0.0 | 1.0 | 0.3 | 7.8 | 0.5 | 0.0 | 9.8 | 0.2 |
| | Mixed species | Mekhlot | 0.3 | 0.0 | 0.0 | 1.0 | 0.3 | 7.8 | 0.5 | 0.0 | 9.8 | 0.2 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 70.1 | 0.0 | 0.0 | 0.0 | 70.1 | 1.3 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 70.1 | 0.0 | 0.0 | 0.0 | 70.1 | 1.3 |
| Myliobatidae etc. | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.6 | 0.0 |
| | Myliobatidea et al. | Lugmah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.6 | 0.0 |
| Nemipteridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.6 | 0.0 | 0.0 | 28.6 | 0.5 |
| | Scolopsis taeniatus | Ebzimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.6 | 0.0 | 0.0 | 28.6 | 0.5 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Pomacanthidae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Pomacanthus maculosus | Anfooz | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.2 | 0.0 |
| Portunidae | | | 0.0 | 0.7 | 0.0 | 0.0 | 102.4 | 20.1 | 0.0 | 0.0 | 123.3 | 2.3 |
| | Portunus pelagicus | Qabqoob | 0.0 | 0.7 | 0.0 | 0.0 | 102.4 | 20.1 | 0.0 | 0.0 | 123.3 | 2.3 |
| Rachycentridae | | | 0.6 | 1.3 | 0.0 | 0.1 | 0.0 | 24.6 | 0.2 | 0.0 | 26.8 | 0.5 |
| | Rachycentron canadum | Sikkil | 0.6 | 1.3 | 0.0 | 0.1 | 0.0 | 24.6 | 0.2 | 0.0 | 26.8 | 0.5 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 1.6 | 0.3 | 3.3 | 1.6 | 0.0 | 6.8 | 0.1 |
| | Scarus ghobban | Gain | 0.0 | 0.0 | 0.0 | 1.6 | 0.3 | 3.3 | 1.6 | 0.0 | 6.8 | 0.1 |

Table 2 - Total catch (tonnes) by landing site in the Emirate of Abu Dhabi (cont.).

| Family | Species | Arabic name | Sila | Delma | Jebel Dhanna | Marfa | Al Bateen | Free Port | Al Sadar | Ghantoot | Total | (%) |
|-----------------------|---------------------------|----------------|-------------|--------------|--------------|--------------|-------------|--------------|--------------|------------|----------------|-------------|
| Scombridae | | | 74.7 | 42.3 | 0.0 | 47.3 | 68.1 | 64.8 | 136.6 | 0.0 | 433.7 | 8.1 |
| | Euthynnus affinis | Sadah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.7 | 0.0 |
| | Scomberomorus commerson | Kanaad | 74.7 | 41.7 | 0.0 | 47.3 | 68.1 | 64.6 | 136.6 | 0.0 | 432.9 | 8.1 |
| Scyllaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Thenus orientalis | Umm Ar Rubiyan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.0 |
| Sepiidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Sepia pharaonis | Habbar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.0 |
| Serranidae | | | 31.4 | 239.6 | 0.0 | 108.4 | 4.5 | 820.4 | 60.3 | 0.0 | 1,264.6 | 23.7 |
| | Cephalopholis hemistiktos | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Epinephelus areolatus | Semman | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Epinephelus coioides | Hamour | 31.4 | 239.5 | 0.0 | 108.4 | 4.5 | 819.8 | 60.3 | 0.0 | 1,264.1 | 23.7 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 15.5 | 15.8 | 0.0 | 0.0 | 31.3 | 0.6 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 15.5 | 15.8 | 0.0 | 0.0 | 31.3 | 0.6 |
| | Siganus javus | Safi Sneifi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sparidae | | | 0.1 | 11.6 | 0.0 | 3.0 | 28.2 | 191.9 | 3.5 | 0.0 | 238.4 | 4.5 |
| | Acanthopagrus bifasciatus | Faskar | 0.0 | 9.9 | 0.0 | 0.4 | 0.0 | 48.4 | 0.3 | 0.0 | 59.1 | 1.1 |
| | Acanthopagrus latus | Shaam | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 5.2 | 0.0 | 0.0 | 5.4 | 0.1 |
| | Argyrops spinifer | Kofar | 0.0 | 0.1 | 0.0 | 0.2 | 0.0 | 45.8 | 0.2 | 0.0 | 46.4 | 0.9 |
| | Rhabdosargus sarba | Qabit | 0.0 | 1.4 | 0.0 | 2.4 | 28.2 | 92.5 | 3.0 | 0.0 | 127.5 | 2.4 |
| Sphyraenidae | | | 1.7 | 2.7 | 0.0 | 38.7 | 7.3 | 11.7 | 19.9 | 0.0 | 82.0 | 1.5 |
| | Sphyraena jello | Jedd | 1.7 | 2.7 | 0.0 | 38.7 | 7.3 | 11.7 | 19.9 | 0.0 | 82.0 | 1.5 |
| Stromateidae | | | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Pampus argenteus | Zubaidi | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.4 | 0.0 |
| Teraponidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Terapon jarbua | Yelly | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tetraodontidae | | | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Arothron stellatus | Faqel | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Total | | | 218.4 | 906.6 | 0.0 | 481.5 | 471.1 | 2,870.8 | 388.3 | 0.0 | 5,336.6 | 100.0 |
| (%) | | | 4.1 | 17.0 | 0.0 | 9.0 | 8.8 | 53.8 | 7.3 | 0.0 | 100.0 | |

Table 3 - Total catch (tonnes) by boat/gear type in the Emirate of Abu Dhabi.

| Family | Species | Arabic name | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|-----------------------|--------------------------|-------------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|--------------|-------------|
| Ariidae | | | 10.1 | 0.0 | 0.0 | 0.0 | 18.7 | 0.0 | 0.1 | 0.0 | 28.9 | 0.5 |
| | Arius thalassinus | Khan | 10.1 | 0.0 | 0.0 | 0.0 | 18.7 | 0.0 | 0.1 | 0.0 | 28.9 | 0.5 |
| Belonidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| | Belonidae et al. | Haqool | 0.2 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| Carangidae | | | 413.6 | 0.0 | 6.6 | 0.4 | 445.3 | 1.2 | 16.8 | 13.2 | 896.9 | 16.8 |
| | Alepes vari | Gamlouh | 1.6 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 1.8 | 0.0 |
| | Atule mate | Durduman | 125.2 | 0.0 | 0.0 | 0.0 | 57.0 | 0.0 | 3.5 | 0.0 | 185.7 | 3.5 |
| | Carangidae all | Jesh unidentified | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Carangoides bajad | Jesh Um Al | 41.1 | 0.0 | 0.0 | 0.2 | 157.5 | 0.1 | 0.7 | 8.2 | 207.8 | 3.9 |
| | Carangoides | Jesh Zeria | 26.3 | 0.0 | 0.0 | 0.0 | 148.7 | 0.0 | 2.9 | 0.6 | 178.5 | 3.3 |
| | Carangoides | Jesh Sal | 9.3 | 0.0 | 0.0 | 0.0 | 21.2 | 0.0 | 0.0 | 0.0 | 30.5 | 0.6 |
| | Caranx heberi | Jesh Farow | 3.9 | 0.0 | 0.0 | 0.0 | 5.1 | 0.0 | 0.0 | 0.0 | 9.1 | 0.2 |
| | Caranx ignobilis | Gufdar | 31.1 | 0.0 | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 33.0 | 0.6 |
| | Gnathanodon | Zuraidi | 143.5 | 0.0 | 0.0 | 0.0 | 12.6 | 0.0 | 0.0 | 0.0 | 156.0 | 2.9 |
| | Megalaspis cordyla | Diyayuh | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 |
| | Parastromateus niger | Halwayoh | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 |
| | Scomberoides | Dhil'e | 9.0 | 0.0 | 0.0 | 0.1 | 9.4 | 1.1 | 9.0 | 3.7 | 32.2 | 0.6 |
| | Scomberoides tol | Bassar | 4.3 | 0.0 | 6.6 | 0.1 | 26.0 | 0.0 | 0.7 | 0.6 | 38.2 | 0.7 |
| | Seriola dumerili | Jib | 13.6 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 19.1 | 0.4 |
| | Seriolina nigrofasciata | Hamam | 3.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.6 | 0.1 |
| Carcharhinidae | | | 0.8 | 0.0 | 7.7 | 0.0 | 4.1 | 0.0 | 0.6 | 0.0 | 13.2 | 0.2 |
| | Carcharhinidae et al. | Jarjoor | 0.8 | 0.0 | 7.7 | 0.0 | 4.1 | 0.0 | 0.6 | 0.0 | 13.2 | 0.2 |
| Chanidae | | | 0.1 | 0.0 | 4.7 | 2.5 | 1.7 | 1.3 | 0.2 | 0.0 | 10.6 | 0.2 |
| | Chanos chanos | Aifah | 0.1 | 0.0 | 4.7 | 2.5 | 1.7 | 1.3 | 0.2 | 0.0 | 10.6 | 0.2 |
| Clupeidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Sardinella longiceps | Oumah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ephippidae | | | 22.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 22.5 | 0.4 |
| | Ephippidae et al. | Imad | 22.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 22.5 | 0.4 |
| Gerreidae | | | 0.2 | 0.0 | 35.6 | 4.0 | 0.1 | 6.4 | 0.0 | 11.8 | 58.2 | 1.1 |
| | Gerres longirostris | Badah | 0.2 | 0.0 | 35.6 | 4.0 | 0.1 | 6.4 | 0.0 | 11.8 | 58.2 | 1.1 |
| Haemulidae | | | 657.4 | 0.0 | 22.2 | 1.4 | 36.1 | 0.7 | 1.8 | 0.0 | 719.7 | 13.5 |
| | Diagramma pictum | Farsh | 589.6 | 0.0 | 0.0 | 0.0 | 32.0 | 0.0 | 1.8 | 0.0 | 623.4 | 11.7 |
| | Plectorhinchus gaterinus | Hilali | 24.7 | 0.0 | 0.0 | 0.0 | 1.4 | 0.0 | 0.0 | 0.0 | 26.1 | 0.5 |
| | Plectorhinchus | Yanam | 42.7 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.1 | 0.0 | 43.2 | 0.8 |
| | Pomadasys argenteus | Naqroor | 0.4 | 0.0 | 22.2 | 1.4 | 2.3 | 0.7 | 0.0 | 0.0 | 27.0 | 0.5 |

Table 3 - Total catch (tonnes) by boat/gear type in the Emirate of Abu Dhabi (cont.).

| Family | Species | Arabic name | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|--------------------------|-----------------------|-----------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|----------------|-------------|
| Labridae | | | 0.6 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 2.0 | 0.0 | 4.3 | 0.1 |
| | Choerodon robustus | Jin | 0.6 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 2.0 | 0.0 | 4.3 | 0.1 |
| Lethrinidae | | | 737.0 | 0.0 | 0.0 | 0.0 | 454.9 | 0.1 | 1.6 | 6.6 | 1,200.3 | 22.5 |
| | Lethrinus borbonicus | Yemah | 51.6 | 0.0 | 0.0 | 0.0 | 10.2 | 0.0 | 0.1 | 0.0 | 61.8 | 1.2 |
| | Lethrinus lentjan | Shaari Eshkheli | 118.1 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 121.0 | 2.3 |
| | Lethrinus microdon | Souli | 30.2 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 35.7 | 0.7 |
| | Lethrinus nebulosus | Shaari | 537.2 | 0.0 | 0.0 | 0.0 | 436.3 | 0.1 | 1.5 | 6.6 | 981.7 | 18.4 |
| Lutjanidae | | | 9.4 | 0.0 | 0.7 | 38.1 | 14.9 | 0.0 | 0.0 | 0.0 | 63.1 | 1.2 |
| | Lutjanus ehrenbergii | Naiser | 2.3 | 0.0 | 0.7 | 38.1 | 9.0 | 0.0 | 0.0 | 0.0 | 50.1 | 0.9 |
| | Lutjanus fulviflamma | Aqalah | 5.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.3 | 0.1 |
| | Lutjanus malabaricus | Hamra | 1.4 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 2.1 | 0.0 |
| | Lutjanus russelli | Umm Dhrais | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Pinjalo pinjalo | Neimi | 0.0 | 0.0 | 0.0 | 0.0 | 5.1 | 0.0 | 0.0 | 0.0 | 5.1 | 0.1 |
| Mixed | | | 6.8 | 0.0 | 0.0 | 0.3 | 2.7 | 0.0 | 0.0 | 0.0 | 9.8 | 0.2 |
| | Mixed species | Mekhlot | 6.8 | 0.0 | 0.0 | 0.3 | 2.7 | 0.0 | 0.0 | 0.0 | 9.8 | 0.2 |
| Mugilidae | | | 0.0 | 0.0 | 62.9 | 0.2 | 0.0 | 0.6 | 0.0 | 6.4 | 70.1 | 1.3 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 62.9 | 0.2 | 0.0 | 0.6 | 0.0 | 6.4 | 70.1 | 1.3 |
| Myliobatidae etc. | | | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 |
| | Myliobatidea et al. | Lugmah | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 |
| Nemipteridae | | | 28.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.6 | 0.5 |
| | Scolopsis taeniatus | Ebzimi | 28.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.6 | 0.5 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.3 | 0.4 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.3 | 0.4 | 0.0 |
| Pomacanthidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Pomacanthus | Anfooz | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Portunidae | | | 20.8 | 0.0 | 51.8 | 3.5 | 0.0 | 0.5 | 0.0 | 46.6 | 123.3 | 2.3 |
| | Portunus pelagicus | Qabqoob | 20.8 | 0.0 | 51.8 | 3.5 | 0.0 | 0.5 | 0.0 | 46.6 | 123.3 | 2.3 |
| Rachycentridae | | | 25.5 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 26.8 | 0.5 |
| | Rachycentron canadum | Sikkil | 25.5 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 26.8 | 0.5 |
| Scaridae | | | 3.3 | 0.0 | 0.0 | 0.0 | 3.2 | 0.3 | 0.0 | 0.0 | 6.8 | 0.1 |
| | Scarus ghobban | Gain | 3.3 | 0.0 | 0.0 | 0.0 | 3.2 | 0.3 | 0.0 | 0.0 | 6.8 | 0.1 |

Table 3 - Total catch (tonnes) by boat/gear type in the Emirate of Abu Dhabi (cont.).

| Family | Species | Arabic name | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|-----------------------|------------------------|-------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|----------------|-------------|
| Scombridae | | | 12.3 | 0.0 | 0.0 | 5.1 | 237.3 | 0.9 | 173.5 | 4.6 | 433.7 | 8.1 |
| | Euthynnus affinis | Sadah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Rastrelliger kanagurta | Garfah | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 |
| | Scomberomorus | Kanaad | 11.5 | 0.0 | 0.0 | 5.1 | 237.3 | 0.9 | 173.5 | 4.6 | 432.9 | 8.1 |
| Scyllaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Thenus orientalis | Umm Ar | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Sepiidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Sepia pharaonis | Habbar | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Serranidae | | | 923.3 | 0.0 | 0.0 | 0.0 | 335.8 | 0.0 | 2.5 | 3.0 | 1,264.6 | 23.7 |
| | Cephalopholis | Eshnenuh | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Epinephelus areolatus | Semman | 0.2 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Epinephelus coioides | Hamour | 923.0 | 0.0 | 0.0 | 0.0 | 335.6 | 0.0 | 2.5 | 3.0 | 1,264.1 | 23.7 |
| Siganidae | | | 15.8 | 0.0 | 0.5 | 1.7 | 0.0 | 2.9 | 0.0 | 10.5 | 31.3 | 0.6 |
| | Siganus canaliculatus | Safi Arabi | 15.8 | 0.0 | 0.5 | 1.7 | 0.0 | 2.9 | 0.0 | 10.5 | 31.3 | 0.6 |
| | Siganus javus | Safi Sneifi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sparidae | | | 137.7 | 0.0 | 1.6 | 1.3 | 96.5 | 0.9 | 0.0 | 0.3 | 238.4 | 4.5 |
| | Acanthopagrus | Faskar | 57.3 | 0.0 | 0.0 | 0.0 | 1.8 | 0.0 | 0.0 | 0.0 | 59.1 | 1.1 |
| | Acanthopagrus latus | Shaam | 4.8 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 5.4 | 0.1 |
| | Argyrops spinifer | Kofar | 45.9 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 46.4 | 0.9 |
| | Rhabdosargus sarba | Qabit | 29.7 | 0.0 | 1.6 | 1.3 | 93.7 | 0.9 | 0.0 | 0.3 | 127.5 | 2.4 |
| Sphyraenidae | | | 1.7 | 0.0 | 0.0 | 0.0 | 72.0 | 0.0 | 5.4 | 2.9 | 82.0 | 1.5 |
| | Sphyraena jello | Jedd | 1.7 | 0.0 | 0.0 | 0.0 | 72.0 | 0.0 | 5.4 | 2.9 | 82.0 | 1.5 |
| Stromateidae | | | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Pampus argenteus | Zubaidi | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Teraponidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Terapon jarbua | Yelly | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tetraodontidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Arothron stellatus | Faqel | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Total | | | 3,029.2 | 0.0 | 194.3 | 58.6 | 1,727.8 | 16.0 | 204.7 | 106.0 | 5,336.6 | 100.0 |
| (%) | | | 56.8 | 0.0 | 3.6 | 1.1 | 32.4 | 0.3 | 3.8 | 2.0 | 100.0 | |

Table 4 (a) - Total catch (tonnes) and species composition for landings at Sila.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-------------------|---------------------------|-----------------|------------|------------|------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 0.0 | 0.0 | 13.2 | 12.8 | 0.0 | 6.0 | 0.0 | 4.1 | 0.0 | 3.0 | 0.0 | 3.6 | 42.8 | 19.6 |
| Carangidae | | | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.3 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Carangoides malabaricus | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Caranx ignobilis | Gufdar | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Gnathanodon speciosus | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Haemulidae | | | 0.0 | 0.0 | 9.7 | 11.2 | 0.0 | 5.2 | 0.0 | 4.1 | 0.0 | 2.5 | 0.0 | 3.6 | 36.3 | 16.6 |
| | Diagramma pictum | Farsh | 0.0 | 0.0 | 9.7 | 11.2 | 0.0 | 5.2 | 0.0 | 4.1 | 0.0 | 2.3 | 0.0 | 3.6 | 36.1 | 16.5 |
| | Plectorhinchus sordidus | Yanam | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.2 | 0.1 |
| Serranidae | | | 0.0 | 0.0 | 3.4 | 1.6 | 0.0 | 0.3 | 0.0 | 0.1 | 0.0 | 0.5 | 0.0 | 0.0 | 5.8 | 2.7 |
| | Epinephelus coioides | Hamour | 0.0 | 0.0 | 3.4 | 1.6 | 0.0 | 0.3 | 0.0 | 0.1 | 0.0 | 0.5 | 0.0 | 0.0 | 5.8 | 2.7 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Acanthopagrus bifasciatus | Faskar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

Table 4 (a) - Total catch (tonnes) and species composition for landings at Sila (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|-------------------------|-------------------|-------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|------------|-------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 39.5 | 26.6 | 12.8 | 9.1 | 5.7 | 11.0 | 4.8 | 6.3 | 4.6 | 9.0 | 18.4 | 11.0 | 158.9 | 72.7 |
| Carangidae | | | 0.4 | 0.9 | 1.6 | 4.0 | 4.5 | 3.2 | 0.6 | 0.5 | 2.3 | 2.0 | 10.7 | 1.5 | 32.4 | 14.8 |
| | Atule mate | Durduman | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Carangidae all | Jesh unidentified | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Carangoides bajad | Jesh Um Al Hala | 0.1 | 0.5 | 0.8 | 1.9 | 3.7 | 1.7 | 0.4 | 0.5 | 2.3 | 1.6 | 10.6 | 1.4 | 25.5 | 11.7 |
| | Carangoides malabaricus | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.4 |
| | Caranx ignobilis | Gufdar | 0.1 | 0.1 | 0.0 | 0.0 | 0.1 | 0.2 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.8 | 0.4 |
| | Gnathanodon speciosus | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.2 |
| | Scomberoides tol | Bassar | 0.0 | 0.3 | 0.8 | 2.1 | 0.5 | 0.0 | 0.2 | 0.0 | 0.0 | 0.1 | 0.2 | 0.0 | 4.3 | 1.9 |
| Carcharhinidae | | | 0.0 | 1.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.5 | 0.7 |
| | Carcharhinidae et al. | Jarjoor | 0.0 | 1.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.5 | 0.7 |
| Chanidae | | | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.8 |
| | Chanos chanos | Aifah | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.8 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| Haemulidae | | | 0.0 | 0.9 | 0.5 | 2.1 | 0.2 | 5.3 | 0.6 | 3.1 | 1.4 | 2.5 | 0.0 | 5.3 | 21.9 | 10.0 |
| | Diagramma pictum | Farsh | 0.0 | 0.9 | 0.5 | 2.1 | 0.2 | 5.3 | 0.6 | 3.1 | 1.4 | 2.5 | 0.0 | 5.3 | 21.9 | 10.0 |
| Lethrinidae | | | 0.3 | 2.6 | 0.6 | 1.5 | 0.4 | 0.2 | 0.9 | 0.0 | 0.0 | 0.4 | 1.8 | 2.0 | 10.5 | 4.8 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.2 |
| | Lethrinus microdon | Souli | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.3 | 0.1 |
| | Lethrinus nebulosus | Shaari | 0.1 | 2.5 | 0.6 | 1.0 | 0.4 | 0.2 | 0.9 | 0.0 | 0.0 | 0.4 | 1.8 | 1.9 | 9.8 | 4.5 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| Rachycentridae | | | 0.1 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.3 |
| | Rachycentron canadum | Sikkil | 0.1 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.3 |
| Scombridae | | | 36.4 | 19.1 | 9.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 65.0 | 29.7 |
| | Scomberomorus commerson | Kanaad | 36.4 | 19.1 | 9.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 65.0 | 29.7 |
| Serranidae | | | 0.4 | 0.4 | 0.5 | 1.1 | 0.5 | 2.4 | 2.7 | 2.6 | 0.8 | 4.1 | 5.8 | 2.1 | 23.5 | 10.8 |
| | Epinephelus coioides | Hamour | 0.4 | 0.4 | 0.5 | 1.1 | 0.5 | 2.4 | 2.7 | 2.6 | 0.8 | 4.1 | 5.8 | 2.1 | 23.5 | 10.8 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sphyraenidae | | | 0.3 | 1.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.7 |
| | Sphyraena jello | Jedd | 0.3 | 1.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.7 |

Table 4 (a) - Total catch (tonnes) and species composition for landings at Sila (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|--------------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Halaq</i> | | 7.5 | 6.0 | 3.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 16.8 | 7.7 |
| Ariidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.8 | 0.2 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.6 |
| | <i>Atule mate</i> | Durduman | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.3 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.3 |
| | <i>Caranx ignobilis</i> | Gufdar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Scomberoides tol</i> | Bassar | 0.5 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.3 |
| Carcharhinidae | | | 0.0 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.2 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.2 |
| Chanidae | | | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | <i>Chanos chanos</i> | Aifah | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| Haemulidae | | | 0.7 | 1.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.8 | 0.8 |
| | <i>Diagramma pictum</i> | Farsh | 0.7 | 1.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.8 | 0.8 |
| | <i>Plectorhinchus sordidus</i> | Yanam | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Lethrinidae | | | 0.1 | 0.6 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.4 |
| | <i>Lethrinus nebulosus</i> | Shaari | 0.1 | 0.6 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.4 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Scombridae | | | 4.2 | 2.8 | 2.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.7 | 4.4 |
| | <i>Scomberomorus commerson</i> | Kanaad | 4.2 | 2.8 | 2.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.7 | 4.4 |
| Serranidae | | | 1.4 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.2 | 1.0 |
| | <i>Epinephelus coioides</i> | Hamour | 1.4 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.2 | 1.0 |
| Sphyraenidae | | | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | <i>Sphyraena jello</i> | Jedd | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| Total | | | 47.0 | 32.6 | 29.4 | 21.9 | 5.7 | 17.0 | 4.8 | 10.4 | 4.6 | 12.0 | 18.4 | 14.7 | 218.4 | 100.0 |
| (%) | | | 21.5 | 14.9 | 13.5 | 10.0 | 2.6 | 7.8 | 2.2 | 4.8 | 2.1 | 5.5 | 8.4 | 6.7 | 100.0 | |

Table 4 (b) - Total catch (tonnes) and species composition for landings at Delma Island.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|----------------------|--------------------------|-----------------|-------------|------------|------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|--------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 45.8 | 20.5 | 30.6 | 120.4 | 82.6 | 109.0 | 78.5 | 36.1 | 48.0 | 51.4 | 55.2 | 68.2 | 746.3 | 82.3 |
| Ariidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.5 | 0.1 |
| | Arius thalassinus | Khan | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.5 | 0.1 |
| Carangidae | | | 4.3 | 2.1 | 1.0 | 3.8 | 9.4 | 20.2 | 6.8 | 2.7 | 2.2 | 3.5 | 3.2 | 1.8 | 61.0 | 6.7 |
| | Alepes vari | Gamlouh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Atule mate | Durduman | 0.0 | 0.0 | 0.0 | 1.0 | 4.8 | 7.7 | 0.8 | 0.3 | 0.1 | 0.8 | 0.1 | 0.1 | 15.8 | 1.7 |
| | Carangoides bajad | Jesh Um Al Hala | 0.4 | 0.0 | 0.0 | 0.0 | 0.3 | 1.2 | 0.4 | 0.2 | 0.1 | 0.3 | 0.1 | 0.0 | 2.9 | 0.3 |
| | Carangoides malabaricus | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Caranx heberi | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Caranx ignobilis | Gufdar | 0.4 | 0.4 | 0.1 | 0.4 | 2.7 | 5.0 | 1.6 | 0.4 | 0.8 | 0.5 | 0.5 | 0.3 | 13.0 | 1.4 |
| | Gnathanodon speciosus | Zuraidi | 3.4 | 1.7 | 1.0 | 2.3 | 0.9 | 4.6 | 3.6 | 1.8 | 1.2 | 1.9 | 2.5 | 1.4 | 26.4 | 2.9 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.2 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.1 |
| | Seriola dumerili | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 1.3 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.2 |
| | Seriolina nigrofasciata | Hamam | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Ephippidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.6 | 2.7 | 0.7 | 0.2 | 0.0 | 5.2 | 0.6 |
| | Ephippidae et al. | Imad | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.6 | 2.7 | 0.7 | 0.2 | 0.0 | 5.2 | 0.6 |
| Haemulidae | | | 3.7 | 0.0 | 0.0 | 65.6 | 32.5 | 39.8 | 34.4 | 18.3 | 23.5 | 23.8 | 26.9 | 34.4 | 303.0 | 33.4 |
| | Diagramma pictum | Farsh | 3.7 | 0.0 | 0.0 | 65.6 | 32.4 | 39.8 | 34.4 | 18.3 | 23.5 | 23.8 | 26.9 | 34.3 | 302.8 | 33.4 |
| | Plectorhinchus gaterinus | Hilali | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Plectorhinchus sordidus | Yanam | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 | 0.0 |
| Labridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Choerodon robustus | Jin | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Lethrinidae | | | 11.0 | 1.3 | 8.4 | 19.5 | 24.3 | 26.0 | 17.8 | 4.7 | 4.0 | 5.3 | 8.3 | 15.0 | 145.8 | 16.1 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.0 | 0.0 | 0.9 | 0.4 | 0.3 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 2.1 | 0.2 |
| | Lethrinus microdon | Souli | 0.0 | 0.0 | 0.0 | 2.5 | 2.0 | 1.9 | 1.3 | 0.5 | 0.3 | 0.5 | 0.6 | 2.0 | 11.6 | 1.3 |
| | Lethrinus nebulosus | Shaari | 11.0 | 1.3 | 8.4 | 16.2 | 21.8 | 23.8 | 16.2 | 4.2 | 3.7 | 4.8 | 7.7 | 12.9 | 132.1 | 14.6 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Lutjanus malabaricus | Hamra | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Lutjanus russelli | Umm Dhrais | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Pomacanthidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Pomacanthus maculosus | Anfooz | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 | 0.4 | 0.7 | 0.1 |
| | Portunus pelagicus | Qabqoob | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 | 0.4 | 0.7 | 0.1 |

Table 4 (b) - Total catch (tonnes) and species composition for landings at Delma Island (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|---------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|------------|-------------|-------------|-------------|-------------|--------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 45.8 | 20.5 | 30.6 | 120.4 | 82.6 | 109.0 | 78.5 | 36.1 | 48.0 | 51.4 | 55.2 | 68.2 | 746.3 | 82.3 |
| Rachycentridae | | | 0.1 | 0.0 | 0.0 | 0.1 | 0.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.9 | 0.1 |
| | Rachycentron canadum | Sikkil | 0.1 | 0.0 | 0.0 | 0.1 | 0.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.9 | 0.1 |
| Scombridae | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.4 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.1 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.1 |
| | Scomberomorus commerson | Kanaad | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Serranidae | | | 23.8 | 16.6 | 20.2 | 29.4 | 14.5 | 21.1 | 18.2 | 9.7 | 15.4 | 17.8 | 16.3 | 15.3 | 218.3 | 24.1 |
| | Epinephelus coioides | Hamour | 23.8 | 16.6 | 20.2 | 29.4 | 14.5 | 21.1 | 18.2 | 9.7 | 15.4 | 17.8 | 16.3 | 15.3 | 218.3 | 24.1 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sparidae | | | 2.4 | 0.5 | 0.9 | 1.9 | 0.8 | 1.1 | 0.4 | 0.1 | 0.2 | 0.1 | 0.1 | 0.9 | 9.3 | 1.0 |
| | Acanthopagrus bifasciatus | Faskar | 2.4 | 0.5 | 0.9 | 1.8 | 0.7 | 1.0 | 0.3 | 0.1 | 0.2 | 0.1 | 0.1 | 0.9 | 9.0 | 1.0 |
| | Acanthopagrus latus | Shaam | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Argyrops spinifer | Kofar | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Sphyraenidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Sphyraena jello | Jedd | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Stromateidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Pampus argenteus | Zubaidi | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Teraponidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Terapon jarbua | Yelly | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tetraodontidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Arothron stellatus | Faqel | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |

Table 4 (b) - Total catch (tonnes) and species composition for landings at Delma Island (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|----------------------------------|-----------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|-------------|-------------|-------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 40.4 | 4.0 | 18.0 | 9.3 | 6.3 | 2.5 | 5.1 | 2.6 | 5.6 | 21.5 | 20.2 | 24.8 | 160.3 | 17.7 |
| Ariidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.4 | 0.0 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.4 | 0.0 |
| Belonidae | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Belonidae et al.</i> | Haqool | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Carangidae | | | 5.4 | 0.0 | 2.9 | 5.4 | 3.9 | 0.5 | 2.4 | 0.8 | 3.2 | 3.7 | 1.9 | 0.8 | 30.9 | 3.4 |
| | <i>Atule mate</i> | Durduman | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 1.3 | 0.1 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 2.8 | 0.0 | 0.0 | 3.7 | 3.5 | 0.2 | 2.3 | 0.4 | 1.4 | 1.1 | 1.3 | 0.8 | 17.6 | 1.9 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Caranx ignobilis</i> | Gufdar | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 1.1 | 0.1 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 1.6 | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 4.7 | 0.5 |
| | <i>Scomberoides</i> | Dhil'e | 0.1 | 0.0 | 0.0 | 0.6 | 0.2 | 0.2 | 0.1 | 0.2 | 1.4 | 2.5 | 0.5 | 0.0 | 5.8 | 0.6 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 |
| Carcharhinidae | | | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.8 | 0.1 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.8 | 0.1 |
| Haemulidae | | | 1.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.8 | 0.2 |
| | <i>Diagramma pictum</i> | Farsh | 1.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.8 | 0.2 |
| Lethrinidae | | | 15.6 | 4.0 | 9.1 | 2.7 | 0.7 | 0.5 | 0.7 | 0.1 | 0.2 | 1.0 | 5.2 | 18.4 | 58.3 | 6.4 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 | 0.0 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.3 | 0.0 | 0.1 | 0.3 | 0.3 | 1.3 | 2.5 | 0.3 |
| | <i>Lethrinus nebulosus</i> | Shaari | 15.6 | 4.0 | 9.1 | 2.7 | 0.6 | 0.4 | 0.4 | 0.0 | 0.1 | 0.7 | 4.8 | 17.1 | 55.6 | 6.1 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Rachycentridae | | | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Scombridae | | | 2.8 | 0.0 | 0.0 | 0.0 | 0.9 | 1.3 | 1.4 | 1.4 | 1.9 | 15.3 | 12.4 | 4.1 | 41.5 | 4.6 |
| | <i>Scomberomorus commerson</i> | Kanaad | 2.8 | 0.0 | 0.0 | 0.0 | 0.9 | 1.3 | 1.4 | 1.4 | 1.9 | 15.3 | 12.4 | 4.1 | 41.5 | 4.6 |
| Serranidae | | | 12.6 | 0.0 | 6.0 | 0.4 | 0.2 | 0.1 | 0.4 | 0.0 | 0.2 | 0.3 | 0.4 | 0.7 | 21.3 | 2.3 |
| | <i>Cephalopholis hemistiktos</i> | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Epinephelus coioides</i> | Hamour | 12.6 | 0.0 | 6.0 | 0.4 | 0.2 | 0.1 | 0.4 | 0.0 | 0.2 | 0.3 | 0.4 | 0.7 | 21.2 | 2.3 |
| Sparidae | | | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 1.1 | 0.0 | 0.0 | 2.4 | 0.3 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 0.1 |
| | <i>Rhabdosargus sarba</i> | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 1.1 | 0.0 | 0.0 | 1.4 | 0.2 |

Table 4 (b) - Total catch (tonnes) and species composition for landings at Delma Island (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|---------------------|------------------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 40.4 | 4.0 | 18.0 | 9.3 | 6.3 | 2.5 | 5.1 | 2.6 | 5.6 | 21.5 | 20.2 | 24.8 | 160.3 | 17.7 |
| Sphyraenidae | | | 0.2 | 0.0 | 0.0 | 0.8 | 0.7 | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | 0.2 | 0.3 | 2.6 | 0.3 |
| | <i>Sphyraena jello</i> | Jedd | 0.2 | 0.0 | 0.0 | 0.8 | 0.7 | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | 0.2 | 0.3 | 2.6 | 0.3 |
| Total | | | 86.2 | 24.5 | 48.6 | 129.7 | 88.9 | 111.5 | 83.6 | 38.7 | 53.7 | 73.0 | 75.4 | 92.9 | 906.6 | 100.0 |
| (%) | | | 9.5 | 2.7 | 5.4 | 14.3 | 9.8 | 12.3 | 9.2 | 4.3 | 5.9 | 8.0 | 8.3 | 10.3 | 100.0 | |

Table 4 (c) - Total catch (tonnes) and species composition for landings at Al Marfa.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|----------------------------------|-----------------|-------------|-------------|-------------|-------------|------------|------------|-------------|------------|------------|------------|------------|-------------|--------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 128.7 | 98.6 | 43.2 | 50.1 | 13.1 | 22.5 | 21.2 | 13.8 | 20.1 | 15.9 | 25.7 | 28.5 | 481.5 | 100.0 |
| Ariidae | | | 1.5 | 2.3 | 0.0 | 2.3 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 6.5 | 1.4 |
| | <i>Arius thalassinus</i> | Khan | 1.5 | 2.3 | 0.0 | 2.3 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 6.5 | 1.4 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | <i>Belonidae et al.</i> | Haqool | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| Carangidae | | | 47.1 | 36.1 | 15.8 | 12.3 | 3.6 | 4.3 | 10.5 | 4.3 | 0.5 | 5.2 | 5.7 | 5.0 | 150.4 | 31.2 |
| | <i>Atule mate</i> | Durduman | 5.9 | 9.4 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 16.3 | 3.4 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.9 | 0.0 | 0.6 | 12.1 | 0.4 | 3.6 | 7.4 | 3.9 | 0.2 | 1.4 | 0.0 | 0.7 | 31.2 | 6.5 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 40.3 | 26.8 | 15.2 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.4 | 5.2 | 4.3 | 98.3 | 20.4 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.1 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 2.8 | 0.4 | 0.3 | 0.0 | 0.3 | 0.0 | 3.9 | 0.8 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 0.0 | 1.0 | 0.9 | 1.4 | 0.0 | 0.0 | 0.0 | 5.8 | 1.2 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 0.0 | 1.6 | 0.0 | 0.0 | 0.0 | 0.9 | 1.4 | 0.0 | 0.0 | 0.0 | 3.9 | 0.8 |
| | <i>Plectorhinchus gaterinus</i> | Hilali | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 0.2 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.2 |
| Labridae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.2 |
| | <i>Choerodon robustus</i> | Jin | 0.0 | 0.0 | 0.0 | 0.2 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.2 |
| Lethrinidae | | | 20.6 | 23.5 | 3.8 | 16.7 | 5.4 | 9.1 | 3.6 | 4.8 | 4.6 | 3.6 | 8.8 | 12.8 | 117.4 | 24.4 |
| | <i>Lethrinus borbonicus</i> | Yemah | 2.1 | 2.3 | 0.4 | 0.0 | 0.5 | 0.0 | 0.0 | 0.2 | 0.0 | 0.3 | 0.5 | 0.0 | 6.4 | 1.3 |
| | <i>Lethrinus lentjan</i> | Shaari Eshkheli | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 | 0.2 | 0.7 | 0.2 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.6 | 0.1 |
| | <i>Lethrinus nebulosus</i> | Shaari | 18.6 | 21.1 | 3.4 | 16.7 | 4.5 | 9.1 | 3.6 | 4.7 | 4.6 | 2.5 | 8.1 | 12.7 | 109.6 | 22.8 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.0 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.0 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 0.2 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 0.2 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 1.6 | 0.3 |
| | <i>Scarus ghobban</i> | Gain | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 1.6 | 0.3 |

Table 4 (c) - Total catch (tonnes) and species composition for landings at Al Marfa (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|---------------------|---------------------------|-------------|-------------|-------------|-------------|-------------|------------|------------|------------|------------|-------------|------------|------------|------------|--------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 128.7 | 98.6 | 43.2 | 50.1 | 13.1 | 22.5 | 21.2 | 13.8 | 20.1 | 15.9 | 25.7 | 28.5 | 481.5 | 100.0 |
| Scombridae | | | 20.6 | 0.0 | 13.3 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 13.1 | 0.0 | 0.0 | 0.0 | 47.3 | 9.8 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Scomberomorus commerson | Kanaad | 20.6 | 0.0 | 13.3 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 13.1 | 0.0 | 0.0 | 0.0 | 47.3 | 9.8 |
| Serranidae | | | 25.0 | 25.8 | 5.7 | 14.5 | 2.2 | 8.7 | 4.1 | 3.2 | 0.6 | 4.1 | 8.1 | 6.3 | 108.4 | 22.5 |
| | Epinephelus coioides | Hamour | 25.0 | 25.8 | 5.7 | 14.5 | 2.2 | 8.7 | 4.1 | 3.2 | 0.6 | 4.1 | 8.1 | 6.3 | 108.4 | 22.5 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.7 | 1.3 | 0.4 | 3.0 | 0.6 |
| | Acanthopagrus bifasciatus | Faskar | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 |
| | Argyrops spinifer | Kofar | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 1.3 | 0.4 | 2.4 | 0.5 |
| Sphyraenidae | | | 13.8 | 10.8 | 4.6 | 0.0 | 0.8 | 0.0 | 1.4 | 0.0 | 0.0 | 1.7 | 1.6 | 4.0 | 38.7 | 8.0 |
| | Sphyraena jello | Jedd | 13.8 | 10.8 | 4.6 | 0.0 | 0.8 | 0.0 | 1.4 | 0.0 | 0.0 | 1.7 | 1.6 | 4.0 | 38.7 | 8.0 |
| Total | | | 128.7 | 98.6 | 43.2 | 50.1 | 13.1 | 22.5 | 21.2 | 13.8 | 20.1 | 15.9 | 25.7 | 28.5 | 481.5 | 100.0 |
| (%) | | | 26.7 | 20.5 | 9.0 | 10.4 | 2.7 | 4.7 | 4.4 | 2.9 | 4.2 | 3.3 | 5.3 | 5.9 | 100.0 | |

Table 4 (d) - Total catch (tonnes) and species composition for landings at Al Bateen.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|------------------------------|-------------|-------------|------------|-------------|-------------|------------|------------|------------|------------|------------|------------|-------------|------------|-------------|-------------|
| <i>Tarad</i> | <i>Al Defara</i> | | 12.2 | 3.3 | 27.0 | 36.9 | 2.7 | 0.0 | 0.0 | 7.5 | 0.0 | 10.7 | 94.0 | 0.0 | 194.3 | 41.3 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 6.0 | 0.0 | 6.6 | 1.4 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 6.0 | 0.0 | 6.6 | 1.4 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.7 | 0.0 | 0.0 | 7.7 | 1.6 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.7 | 0.0 | 0.0 | 7.7 | 1.6 |
| Chanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 0.0 | 0.0 | 2.1 | 0.0 | 0.0 | 0.0 | 0.0 | 4.7 | 1.0 |
| | <i>Chanos chanos</i> | Aifah | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 0.0 | 0.0 | 2.1 | 0.0 | 0.0 | 0.0 | 0.0 | 4.7 | 1.0 |
| Gerreidae | | | 12.2 | 3.3 | 13.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 6.0 | 0.0 | 35.6 | 7.6 |
| | <i>Gerres longirostris</i> | Badah | 12.2 | 3.3 | 13.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 6.0 | 0.0 | 35.6 | 7.6 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 22.2 | 0.0 | 22.2 | 4.7 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 22.2 | 0.0 | 22.2 | 4.7 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 |
| Mugilidae | | | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 59.8 | 0.0 | 62.9 | 13.3 |
| | <i>Valamugil seheli</i> | Beyah Arabi | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 59.8 | 0.0 | 62.9 | 13.3 |
| Portunidae | | | 0.0 | 0.0 | 11.9 | 36.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 51.8 | 11.0 |
| | <i>Portunus pelagicus</i> | Qabqoob | 0.0 | 0.0 | 11.9 | 36.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 51.8 | 11.0 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.1 |
| | <i>Siganus canaliculatus</i> | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.1 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.6 | 0.0 | 0.0 | 0.0 | 0.0 | 1.6 | 0.3 |
| | <i>Rhabdosargus sarba</i> | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.6 | 0.0 | 0.0 | 0.0 | 0.0 | 1.6 | 0.3 |

Table 4 (d) - Total catch (tonnes) and species composition for landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-------------------|-------------------------|-----------------|------------|------------|------------|------------|-------------|-------------|------------|------------|------------|------------|------------|------------|-------------|------------|
| <i>Tarad</i> | <i>Al Hadhra</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 11.4 | 11.6 | 11.0 | 13.7 | 10.8 | 0.0 | 0.0 | 0.0 | 58.6 | 12.4 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Belonidae et al. | Haqool | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Scomberoides tol | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Chanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 | 0.5 |
| | Chanos chanos | Aifah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 | 0.5 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.7 | 3.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.9 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.7 | 3.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.9 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 1.4 | 0.3 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 1.4 | 0.3 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 10.0 | 10.1 | 8.7 | 7.4 | 2.0 | 0.0 | 0.0 | 0.0 | 38.1 | 8.1 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 10.0 | 10.1 | 8.7 | 7.4 | 2.0 | 0.0 | 0.0 | 0.0 | 38.1 | 8.1 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.4 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 | 0.7 |
| | Portunus pelagicus | Qabqoob | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.4 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 | 0.7 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.1 | 0.0 | 0.0 | 0.0 | 5.1 | 1.1 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.1 | 0.0 | 0.0 | 0.0 | 5.1 | 1.1 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.4 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.4 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.3 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.3 |

Table 4 (d) - Total catch (tonnes) and species composition for landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|---------------------|-------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|------------|------------|-------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 2.2 | 0.0 | 0.3 | 0.0 | 5.2 | 22.2 | 7.2 | 3.0 | 40.2 | 8.5 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.2 |
| | Atule mate | Durduman | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.1 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 |
| | Plectorhinchus sordidus | Yanam | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 |
| Lethrinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.7 | 0.1 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 |
| | Lethrinus nebulosus | Shaari | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.6 | 0.1 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.2 | 0.0 | 5.1 | 0.0 | 10.4 | 2.2 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.2 | 0.0 | 5.1 | 0.0 | 10.4 | 2.2 |
| Serranidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 1.5 | 0.3 |
| | Epinephelus coioides | Hamour | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 1.5 | 0.3 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 22.2 | 0.0 | 1.9 | 24.1 | 5.1 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 22.2 | 0.0 | 1.9 | 24.1 | 5.1 |
| Sphyraenidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 0.1 | 2.1 | 0.5 |
| | Sphyraena jello | Jedd | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 0.1 | 2.1 | 0.5 |

Table 4 (d) - Total catch (tonnes) and species composition for landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|---------------------|---------------------------|-------------|-------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|-------------|
| <i>Tarad</i> | <i>Halaq</i> | | 19.8 | 30.4 | 5.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 56.0 | 11.9 |
| Carangidae | | | 0.0 | 6.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.6 | 1.4 |
| | Carangoides fulvoguttatus | Jesh Zeria | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.6 |
| | Scomberoides | Dhil'e | 0.0 | 3.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 0.8 |
| Scombridae | | | 19.8 | 23.8 | 3.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 47.1 | 10.0 |
| | Scomberomorus commerson | Kanaad | 19.8 | 23.8 | 3.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 47.1 | 10.0 |
| Sphyraenidae | | | 0.0 | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.3 | 0.5 |
| | Sphyraena jello | Jedd | 0.0 | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.3 | 0.5 |

Table 4 (d) - Total catch (tonnes) and species composition for landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|------------------------|---------------------------|-----------------|-------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|------------|
| <i>Tarad</i> | <i>Leikh</i> | | 11.6 | 62.6 | 7.5 | 0.0 | 0.0 | 6.1 | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 16.4 | 106.0 | 22.5 |
| Carangidae | | | 0.0 | 11.9 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 13.2 | 2.8 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 8.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.2 | 1.7 |
| | Carangoides fulvoguttatus | Jesh Zeria | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.1 |
| | Scomberoides | Dhil'e | 0.0 | 3.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 0.8 |
| | Scomberoides tol | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.6 | 0.1 |
| Gerreidae | | | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 8.0 | 11.8 | 2.5 |
| | Gerres longirostris | Badah | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 8.0 | 11.8 | 2.5 |
| Lethrinidae | | | 0.0 | 6.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.6 | 1.4 |
| | Lethrinus nebulosus | Shaari | 0.0 | 6.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.6 | 1.4 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.1 | 6.4 | 1.3 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.1 | 6.4 | 1.3 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.1 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.1 |
| Portunidae | | | 11.6 | 23.6 | 7.5 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.3 | 46.6 | 9.9 |
| | Portunus pelagicus | Qabqoob | 11.6 | 23.6 | 7.5 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.3 | 46.6 | 9.9 |
| Scomberidae | | | 0.0 | 2.1 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 4.6 | 1.0 |
| | Scomberomorus commerson | Kanaad | 0.0 | 2.1 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 4.6 | 1.0 |
| Serranidae | | | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 3.0 | 0.6 |
| | Epinephelus coioides | Hamour | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 3.0 | 0.6 |
| Siganidae | | | 0.0 | 9.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 10.5 | 2.2 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 9.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 10.5 | 2.2 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.1 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.1 |
| Sphyraenidae | | | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.6 |
| | Sphyraena jello | Jedd | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.6 |

Table 4 (d) - Total catch (tonnes) and species composition for landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|------------------------|-------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Al Sakkar</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.8 | 2.6 | 0.0 | 0.0 | 0.0 | 7.5 | 16.0 | 3.4 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Belonidae et al. | Haqool | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.3 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 1.1 | 0.2 |
| Chanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.3 |
| | Chanos chanos | Aifah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.3 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 2.2 | 0.0 | 0.0 | 0.0 | 3.1 | 6.4 | 1.4 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.0 | 2.2 | 0.0 | 0.0 | 0.0 | 3.1 | 6.4 | 1.4 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 |
| Lethrinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Lethrinus nebulosus | Shaari | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.6 | 0.1 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.6 | 0.1 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.5 | 0.1 |
| | Portunus pelagicus | Qabqoob | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.5 | 0.1 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| | Scarus ghobban | Gain | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.2 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.2 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 2.8 | 2.9 | 0.6 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 2.8 | 2.9 | 0.6 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.9 | 0.2 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.9 | 0.2 |
| Total | | | 43.7 | 96.3 | 40.3 | 36.9 | 16.4 | 17.7 | 17.1 | 23.9 | 17.9 | 32.9 | 101.2 | 26.9 | 471.1 | 100.0 |
| (%) | | | 9.3 | 20.4 | 8.5 | 7.8 | 3.5 | 3.8 | 3.6 | 5.1 | 3.8 | 7.0 | 21.5 | 5.7 | 100.0 | |

Table 4 (e) - Total catch (tonnes) and species composition for landings at Free Port.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|---------------------------|-----------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|--------------|-------------|
| <i>Launch</i> | <i>Gargoor</i> | | 248.1 | 142.7 | 275.2 | 372.1 | 258.9 | 177.3 | 134.6 | 152.3 | 101.8 | 158.8 | 73.7 | 144.6 | 2,240.2 | 78.0 |
| Ariidae | | | 0.1 | 1.0 | 0.2 | 0.7 | 1.9 | 1.6 | 0.4 | 0.6 | 0.0 | 0.2 | 1.2 | 1.5 | 9.6 | 0.3 |
| | Arius thalassinus | Khan | 0.1 | 1.0 | 0.2 | 0.7 | 1.9 | 1.6 | 0.4 | 0.6 | 0.0 | 0.2 | 1.2 | 1.5 | 9.6 | 0.3 |
| Belonidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Belonidae et al. | Haqool | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Carangidae | | | 26.3 | 7.0 | 13.4 | 23.6 | 75.3 | 47.2 | 19.8 | 26.6 | 27.7 | 37.6 | 20.7 | 26.8 | 352.0 | 12.3 |
| | Alepes vari | Gamlouh | 0.1 | 0.0 | 0.0 | 0.1 | 0.2 | 0.2 | 0.4 | 0.1 | 0.0 | 0.3 | 0.1 | 0.1 | 1.5 | 0.1 |
| | Atule mate | Durduman | 2.3 | 0.5 | 0.7 | 5.1 | 29.4 | 16.3 | 4.7 | 9.5 | 10.8 | 20.5 | 6.1 | 3.5 | 109.4 | 3.8 |
| | Carangoides bajad | Jesh Um Al Hala | 3.9 | 0.5 | 2.0 | 2.6 | 2.6 | 4.8 | 3.2 | 2.2 | 5.9 | 3.5 | 2.3 | 4.3 | 37.9 | 1.3 |
| | Carangoides fulvoguttatus | Jesh Zeria | 0.0 | 1.2 | 1.3 | 2.7 | 7.5 | 2.2 | 1.1 | 1.2 | 0.3 | 3.3 | 4.1 | 1.3 | 26.3 | 0.9 |
| | Carangoides malabaricus | Jesh Sal | 2.0 | 0.5 | 0.0 | 2.0 | 2.4 | 0.1 | 0.1 | 0.2 | 0.2 | 0.0 | 0.5 | 1.0 | 9.0 | 0.3 |
| | Caranx heberi | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.5 | 0.9 | 0.2 | 0.2 | 0.1 | 0.3 | 0.6 | 3.9 | 0.1 |
| | Caranx ignobilis | Gufdar | 0.4 | 0.0 | 0.8 | 0.5 | 1.9 | 3.8 | 0.8 | 4.7 | 0.6 | 0.8 | 2.0 | 1.7 | 17.9 | 0.6 |
| | Gnathanodon speciosus | Zuraidi | 16.1 | 3.2 | 7.3 | 8.7 | 21.2 | 12.4 | 7.6 | 6.8 | 9.1 | 8.4 | 4.7 | 11.6 | 117.1 | 4.1 |
| | Megalaspis cordyla | Diyayuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.0 |
| | Parastromateus niger | Halwayoh | 0.0 | 0.1 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.8 | 0.0 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.1 | 0.5 | 4.6 | 1.6 | 0.0 | 0.2 | 0.0 | 0.6 | 0.2 | 0.4 | 8.2 | 0.3 |
| | Scomberoides tol | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 1.5 | 0.5 | 0.5 | 0.6 | 0.1 | 0.0 | 0.5 | 4.3 | 0.2 |
| | Seriola dumerili | Jib | 0.0 | 0.4 | 0.6 | 1.3 | 4.7 | 2.6 | 0.5 | 1.0 | 0.0 | 0.1 | 0.0 | 0.6 | 11.8 | 0.4 |
| | Seriolina nigrofasciata | Hamam | 1.5 | 0.4 | 0.3 | 0.1 | 0.2 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.8 | 3.5 | 0.1 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.4 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 |
| | Carcharhinidae et al. | Jarjoor | 0.0 | 0.0 | 0.0 | 0.4 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 |
| Chanidae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Chanos chanos | Aifah | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Clupeidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Sardinella longiceps | Oumah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ephippidae | | | 0.0 | 0.1 | 0.1 | 0.0 | 0.1 | 1.5 | 4.0 | 3.0 | 3.8 | 3.7 | 0.4 | 0.7 | 17.3 | 0.6 |
| | Ephippidae et al. | Imad | 0.0 | 0.1 | 0.1 | 0.0 | 0.1 | 1.5 | 4.0 | 3.0 | 3.8 | 3.7 | 0.4 | 0.7 | 17.3 | 0.6 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| Haemulidae | | | 35.3 | 25.7 | 48.5 | 43.4 | 30.3 | 19.4 | 17.7 | 23.5 | 12.8 | 22.6 | 12.0 | 26.8 | 318.0 | 11.1 |
| | Diagramma pictum | Farsh | 20.2 | 15.0 | 32.2 | 38.3 | 27.1 | 18.0 | 16.3 | 21.7 | 11.4 | 19.6 | 9.9 | 21.1 | 250.7 | 8.7 |
| | Plectorhinchus gaterinus | Hilali | 6.1 | 1.5 | 3.4 | 0.9 | 1.8 | 1.1 | 0.8 | 1.4 | 0.9 | 2.0 | 1.4 | 3.4 | 24.7 | 0.9 |
| | Plectorhinchus sordidus | Yanam | 9.0 | 9.2 | 12.7 | 4.0 | 1.4 | 0.3 | 0.7 | 0.4 | 0.5 | 1.0 | 0.7 | 2.3 | 42.3 | 1.5 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.2 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |

Table 4 (e) - Total catch (tonnes) and species composition for landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|--------------------------|-----------------------|-----------------|-------------|-------------|-------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|--------------|-------------|
| <i>Launch</i> | <i>Gargoor</i> | | 248.1 | 142.7 | 275.2 | 372.1 | 258.9 | 177.3 | 134.6 | 152.3 | 101.8 | 158.8 | 73.7 | 144.6 | 2,240.2 | 78.0 |
| Labridae | | | 0.0 | 0.3 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.0 |
| | Choerodon robustus | Jin | 0.0 | 0.3 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.0 |
| Lethrinidae | | | 28.5 | 24.1 | 61.2 | 155.3 | 85.7 | 53.7 | 45.6 | 45.5 | 18.2 | 31.6 | 17.8 | 24.1 | 591.3 | 20.6 |
| | Lethrinus borbonicus | Yemah | 3.9 | 3.0 | 5.8 | 14.7 | 2.2 | 5.6 | 3.9 | 3.2 | 2.3 | 2.0 | 0.8 | 2.2 | 49.5 | 1.7 |
| | Lethrinus lentjan | Shaari Eshkheli | 4.5 | 3.1 | 8.6 | 21.0 | 11.7 | 10.9 | 14.9 | 14.1 | 3.5 | 10.7 | 6.8 | 8.3 | 118.1 | 4.1 |
| | Lethrinus microdon | Souli | 1.2 | 2.1 | 1.6 | 1.5 | 1.6 | 1.8 | 2.0 | 2.5 | 1.2 | 1.6 | 0.7 | 0.9 | 18.6 | 0.6 |
| | Lethrinus nebulosus | Shaari | 18.9 | 16.0 | 45.2 | 118.0 | 70.2 | 35.3 | 24.9 | 25.7 | 11.2 | 17.3 | 9.5 | 12.8 | 405.1 | 14.1 |
| Lutjanidae | | | 0.3 | 0.2 | 1.0 | 1.1 | 1.4 | 1.2 | 0.6 | 0.7 | 0.8 | 0.9 | 0.3 | 0.7 | 9.3 | 0.3 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.0 | 0.0 | 0.7 | 0.3 | 0.8 | 0.1 | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 2.2 | 0.1 |
| | Lutjanus fulviflamma | Aqalah | 0.2 | 0.1 | 1.0 | 0.3 | 1.0 | 0.3 | 0.3 | 0.6 | 0.6 | 0.8 | 0.1 | 0.0 | 5.3 | 0.2 |
| | Lutjanus malabaricus | Hamra | 0.1 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.5 | 1.4 | 0.0 |
| | Lutjanus russelli | Umm Dhrais | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.4 | 0.0 |
| Mixed | | | 0.3 | 0.5 | 0.6 | 1.3 | 0.9 | 0.5 | 0.6 | 0.4 | 0.4 | 0.4 | 0.2 | 0.7 | 6.8 | 0.2 |
| | Mixed species | Mekhlot | 0.3 | 0.5 | 0.6 | 1.3 | 0.9 | 0.5 | 0.6 | 0.4 | 0.4 | 0.4 | 0.2 | 0.7 | 6.8 | 0.2 |
| Myliobatidae etc. | | | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 |
| | Myliobatidea et al. | Lugmah | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 |
| Nemipteridae | | | 3.3 | 4.5 | 6.1 | 4.4 | 0.9 | 1.8 | 0.6 | 0.7 | 1.4 | 1.0 | 0.8 | 3.1 | 28.6 | 1.0 |
| | Scolopsis taeniatus | Ebzimi | 3.3 | 4.5 | 6.1 | 4.4 | 0.9 | 1.8 | 0.6 | 0.7 | 1.4 | 1.0 | 0.8 | 3.1 | 28.6 | 1.0 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Pomacanthidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Pomacanthus maculosus | Anfooz | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Portunidae | | | 0.9 | 0.2 | 0.1 | 0.8 | 0.4 | 0.5 | 0.8 | 1.3 | 1.8 | 5.6 | 2.7 | 4.9 | 20.1 | 0.7 |
| | Portunus pelagicus | Qabqoob | 0.9 | 0.2 | 0.1 | 0.8 | 0.4 | 0.5 | 0.8 | 1.3 | 1.8 | 5.6 | 2.7 | 4.9 | 20.1 | 0.7 |
| Rachycentridae | | | 2.0 | 2.7 | 3.0 | 2.2 | 4.6 | 1.5 | 0.6 | 1.3 | 0.1 | 2.7 | 1.3 | 2.7 | 24.6 | 0.9 |
| | Rachycentron canadum | Sikkil | 2.0 | 2.7 | 3.0 | 2.2 | 4.6 | 1.5 | 0.6 | 1.3 | 0.1 | 2.7 | 1.3 | 2.7 | 24.6 | 0.9 |
| Scaridae | | | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 1.1 | 0.3 | 0.2 | 0.5 | 0.0 | 0.1 | 0.6 | 3.3 | 0.1 |
| | Scarus ghobban | Gain | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 1.1 | 0.3 | 0.2 | 0.5 | 0.0 | 0.1 | 0.6 | 3.3 | 0.1 |

Table 4 (e) - Total catch (tonnes) and species composition for landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|---------------------|----------------------------------|-------------|--------------|-------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|--------------|-------------|
| <i>Launch</i> | <i>Gargoor</i> | | 248.1 | 142.7 | 275.2 | 372.1 | 258.9 | 177.3 | 134.6 | 152.3 | 101.8 | 158.8 | 73.7 | 144.6 | 2,240.2 | 78.0 |
| Scombridae | | | 0.0 | 1.3 | 7.8 | 1.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 1.0 | 0.1 | 0.0 | 11.5 | 0.4 |
| | <i>Euthynnus affinis</i> | Sadah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Rastrelliger kanagurta</i> | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.0 | 1.3 | 7.8 | 1.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 1.0 | 0.0 | 0.0 | 11.3 | 0.4 |
| Serranidae | | | 105.1 | 53.9 | 110.5 | 122.1 | 47.3 | 42.9 | 39.9 | 43.8 | 30.5 | 46.3 | 14.3 | 42.5 | 699.2 | 24.4 |
| | <i>Cephalopholis hemistiktos</i> | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Epinephelus areolatus</i> | Semman | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 |
| | <i>Epinephelus coioides</i> | Hamour | 105.1 | 53.9 | 110.4 | 122.0 | 47.3 | 42.8 | 39.8 | 43.8 | 30.5 | 46.3 | 14.3 | 42.5 | 698.8 | 24.3 |
| Siganidae | | | 0.4 | 0.5 | 0.9 | 4.3 | 1.9 | 0.8 | 1.1 | 1.5 | 1.5 | 2.0 | 0.6 | 0.2 | 15.8 | 0.5 |
| | <i>Siganus canaliculatus</i> | Safi Arabi | 0.4 | 0.5 | 0.9 | 4.3 | 1.9 | 0.8 | 1.1 | 1.5 | 1.5 | 2.0 | 0.6 | 0.2 | 15.8 | 0.5 |
| | <i>Siganus javus</i> | Safi Sneifi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sparidae | | | 44.8 | 20.4 | 21.6 | 10.9 | 7.5 | 3.2 | 2.4 | 2.6 | 2.1 | 2.4 | 1.5 | 9.2 | 128.4 | 4.5 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 10.4 | 6.7 | 10.6 | 2.4 | 2.9 | 2.2 | 2.3 | 2.5 | 2.0 | 2.0 | 1.2 | 3.2 | 48.3 | 1.7 |
| | <i>Acanthopagrus latus</i> | Shaam | 0.5 | 0.5 | 1.4 | 0.8 | 0.2 | 0.3 | 0.1 | 0.1 | 0.0 | 0.3 | 0.0 | 0.3 | 4.6 | 0.2 |
| | <i>Argyrops spinifer</i> | Kofar | 14.3 | 7.6 | 9.4 | 7.4 | 4.4 | 0.6 | 0.1 | 0.0 | 0.1 | 0.0 | 0.2 | 1.8 | 45.8 | 1.6 |
| | <i>Rhabdosargus sarba</i> | Qabit | 19.6 | 5.7 | 0.2 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.9 | 29.7 | 1.0 |
| Sphyraenidae | | | 0.0 | 0.1 | 0.0 | 0.2 | 0.1 | 0.3 | 0.0 | 0.3 | 0.1 | 0.6 | 0.0 | 0.0 | 1.5 | 0.1 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 0.1 | 0.0 | 0.2 | 0.1 | 0.3 | 0.0 | 0.3 | 0.1 | 0.6 | 0.0 | 0.0 | 1.5 | 0.1 |
| Stromateidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.2 | 0.0 |
| | <i>Pampus argenteus</i> | Zubaidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.2 | 0.0 |

Table 4 (e) - Total catch (tonnes) and species composition for landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|----------------------------------|-----------------|-------------|-------------|-------------|-------------|------------|-------------|-------------|------------|-------------|------------|-------------|-------------|--------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 110.1 | 45.3 | 94.2 | 39.2 | 23.1 | 27.6 | 29.2 | 23.6 | 66.9 | 85.1 | 48.5 | 37.7 | 630.6 | 22.0 |
| Ariidae | | | 1.1 | 0.4 | 0.0 | 0.7 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 3.5 | 8.5 | 0.3 |
| | <i>Arius thalassinus</i> | Khan | 1.1 | 0.4 | 0.0 | 0.7 | 0.0 | 2.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 3.5 | 8.5 | 0.3 |
| Carangidae | | | 12.9 | 9.9 | 45.9 | 16.5 | 9.1 | 6.9 | 9.6 | 3.5 | 19.8 | 7.8 | 18.4 | 6.2 | 166.5 | 5.8 |
| | <i>Alepes vari</i> | Gamlouh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Atule mate</i> | Durduman | 7.4 | 1.2 | 13.0 | 3.5 | 3.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.3 | 2.1 | 35.6 | 1.2 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 5.6 | 7.8 | 14.7 | 10.1 | 3.8 | 4.9 | 3.9 | 1.4 | 0.1 | 4.9 | 2.5 | 0.4 | 60.1 | 2.1 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 0.2 | 0.0 | 0.0 | 2.2 | 0.5 | 3.9 | 0.0 | 0.0 | 0.0 | 4.6 | 3.7 | 15.0 | 0.5 |
| | <i>Carangoides malabaricus</i> | Jesh Sal | 0.0 | 0.0 | 0.0 | 2.6 | 0.0 | 0.8 | 0.0 | 0.0 | 11.0 | 0.0 | 5.8 | 0.0 | 20.3 | 0.7 |
| | <i>Caranx heberi</i> | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 2.9 | 1.4 | 0.0 | 0.0 | 5.1 | 0.2 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 7.2 | 0.3 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.1 | 1.1 | 1.0 | 0.1 | 0.0 | 2.7 | 0.1 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 11.3 | 0.0 | 0.0 | 0.2 | 1.7 | 1.0 | 0.9 | 0.5 | 0.0 | 0.0 | 15.5 | 0.5 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 3.6 | 0.0 | 0.0 | 0.0 | 4.8 | 0.2 |
| Carcharhinidae | | | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 1.0 | 0.0 | 0.0 | 1.7 | 0.1 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 1.0 | 0.0 | 0.0 | 1.7 | 0.1 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.6 | 1.6 | 0.1 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 1.2 | 0.0 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.5 | 0.0 |
| Lethrinidae | | | 40.3 | 20.7 | 41.9 | 16.7 | 3.8 | 12.9 | 14.0 | 5.7 | 0.2 | 3.6 | 12.8 | 15.3 | 187.9 | 6.5 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.6 | 0.0 |
| | <i>Lethrinus lentjan</i> | Shaari Eshkheli | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 |
| | <i>Lethrinus microdon</i> | Souli | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 |
| | <i>Lethrinus nebulosus</i> | Shaari | 39.5 | 19.9 | 41.9 | 16.7 | 3.8 | 12.9 | 13.0 | 5.5 | 0.0 | 3.6 | 12.7 | 15.3 | 184.7 | 6.4 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.1 | 0.0 | 8.1 | 5.1 | 0.0 | 0.7 | 0.0 | 14.4 | 0.5 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.1 | 0.0 | 8.1 | 0.0 | 0.0 | 0.0 | 0.0 | 8.5 | 0.3 |
| | <i>Lutjanus fulviflamma</i> | Aqalah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lutjanus malabaricus</i> | Hamra | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.7 | 0.0 |
| | <i>Pinjalo pinjalo</i> | Neimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.1 | 0.0 | 0.0 | 0.0 | 5.1 | 0.2 |
| Mixed | | | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 1.0 | 0.0 |
| | <i>Mixed species</i> | Mekhlot | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 1.0 | 0.0 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

Table 4 (e) - Total catch (tonnes) and species composition for landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|---------------------|----------------------------------|----------------|-------------|-------------|------------|------------|------------|------------|------------|------------|-------------|-------------|-------------|------------|--------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 110.1 | 45.3 | 94.2 | 39.2 | 23.1 | 27.6 | 29.2 | 23.6 | 66.9 | 85.1 | 48.5 | 37.7 | 630.6 | 22.0 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 8.6 | 2.2 | 4.0 | 0.0 | 18.1 | 20.4 | 0.0 | 0.0 | 53.3 | 1.9 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 8.6 | 2.2 | 4.0 | 0.0 | 18.1 | 20.4 | 0.0 | 0.0 | 53.3 | 1.9 |
| Scyllaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | <i>Thenus orientalis</i> | Umm Ar Rubiyan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Sepiidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | <i>Sepia pharaonis</i> | Habbar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| Serranidae | | | 54.9 | 14.3 | 5.3 | 3.3 | 1.5 | 0.7 | 0.9 | 1.7 | 1.0 | 15.6 | 12.1 | 9.9 | 121.2 | 4.2 |
| | <i>Cephalopholis hemistiktos</i> | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Epinephelus areolatus</i> | Semman | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Epinephelus coioides</i> | Hamour | 54.9 | 14.3 | 5.3 | 3.3 | 1.4 | 0.6 | 0.9 | 1.7 | 1.0 | 15.6 | 12.1 | 9.9 | 121.0 | 4.2 |
| Sparidae | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 3.8 | 21.7 | 34.7 | 2.2 | 0.6 | 63.5 | 2.2 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Acanthopagrus latus</i> | Shaam | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 |
| | <i>Rhabdosargus sarba</i> | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.8 | 21.7 | 34.7 | 2.2 | 0.6 | 62.9 | 2.2 |
| Sphyraenidae | | | 0.0 | 0.0 | 1.1 | 1.8 | 0.1 | 1.8 | 0.1 | 0.0 | 1.1 | 2.2 | 1.9 | 0.0 | 10.1 | 0.4 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 0.0 | 1.1 | 1.8 | 0.1 | 1.8 | 0.1 | 0.0 | 1.1 | 2.2 | 1.9 | 0.0 | 10.1 | 0.4 |
| Total | | | 358.2 | 188.0 | 369.4 | 411.4 | 282.0 | 204.9 | 163.8 | 175.9 | 168.8 | 243.9 | 122.2 | 182.3 | 2,870.8 | 100.0 |
| (%) | | | 12.5 | 6.6 | 12.9 | 14.3 | 9.8 | 7.1 | 5.7 | 6.1 | 5.9 | 8.5 | 4.3 | 6.4 | 100.0 | |

Table 4 (f) - Total catch (tonnes) and species composition for landings at Al Sadar.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|---------------------------|-----------------|------------|------------|------------|-------------|------------|------------|------------|------------|------------|------------|-------------|-------------|-------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 0.0 | 24.8 | 21.1 | 48.9 | 16.8 | 13.0 | 9.9 | 5.7 | 20.1 | 18.5 | 31.3 | 46.2 | 256.4 | 66.0 |
| Ariidae | | | 0.0 | 0.6 | 0.0 | 2.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 3.4 | 0.9 |
| | Arius thalassinus | Khan | 0.0 | 0.6 | 0.0 | 2.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 3.4 | 0.9 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Belonidae et al. | Haqool | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| Carangidae | | | 0.0 | 9.1 | 7.7 | 12.0 | 4.6 | 2.5 | 4.9 | 1.8 | 0.5 | 6.0 | 6.9 | 8.1 | 64.2 | 16.5 |
| | Atule mate | Durduman | 0.0 | 2.4 | 0.0 | 0.0 | 0.1 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 3.1 | 0.8 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.3 | 11.8 | 0.6 | 2.1 | 3.5 | 1.6 | 0.2 | 1.6 | 0.0 | 1.1 | 22.8 | 5.9 |
| | Carangoides fulvoguttatus | Jesh Zeria | 0.0 | 6.7 | 7.4 | 0.0 | 3.9 | 0.0 | 0.0 | 0.0 | 0.0 | 3.9 | 6.4 | 7.0 | 35.3 | 9.1 |
| | Gnathanodon speciosus | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.5 | 0.1 |
| | Scomberoides tol | Bassar | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 1.3 | 0.2 | 0.3 | 0.0 | 0.3 | 0.0 | 2.3 | 0.6 |
| | Seriola dumerili | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Carcharhinidae et al. | Jarjoor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 0.0 | 0.4 | 0.4 | 1.4 | 0.0 | 0.0 | 0.0 | 4.7 | 1.2 |
| | Diagramma pictum | Farsh | 0.0 | 0.0 | 0.0 | 1.6 | 0.0 | 0.0 | 0.0 | 0.4 | 1.4 | 0.0 | 0.0 | 0.0 | 3.3 | 0.9 |
| | Plectorhinchus gaterinus | Hilali | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.2 |
| Labridae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.2 |
| | Choerodon robustus | Jin | 0.0 | 0.0 | 0.0 | 0.2 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.2 |
| Lethrinidae | | | 0.0 | 5.9 | 1.9 | 16.3 | 6.9 | 5.3 | 1.7 | 2.0 | 4.6 | 4.2 | 10.6 | 20.8 | 80.2 | 20.6 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.6 | 0.2 | 0.0 | 0.7 | 0.0 | 0.0 | 0.1 | 0.0 | 0.4 | 0.6 | 0.0 | 2.6 | 0.7 |
| | Lethrinus lentjan | Shaari Eshkheli | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 | 0.3 | 0.9 | 0.2 |
| | Lethrinus microdon | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.8 | 0.2 |
| | Lethrinus nebulosus | Shaari | 0.0 | 5.3 | 1.7 | 16.3 | 5.7 | 5.3 | 1.7 | 1.9 | 4.6 | 3.0 | 9.9 | 20.5 | 75.9 | 19.5 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.2 | 0.1 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.2 | 0.1 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.1 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.1 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.0 |
| | Rachycentron canadum | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.0 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 1.6 | 0.4 |
| | Scarus ghobban | Gain | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 1.6 | 0.4 |

Table 4 (f) - Total catch (tonnes) and species composition for landings at Al Sadar (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|---------------------|---------------------------|-------------|------------|------------|------------|-------------|------------|------------|------------|------------|-------------|------------|------------|-------------|-------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 0.0 | 24.8 | 21.1 | 48.9 | 16.8 | 13.0 | 9.9 | 5.7 | 20.1 | 18.5 | 31.3 | 46.2 | 256.4 | 66.0 |
| Scombridae | | | 0.0 | 0.0 | 6.5 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 13.1 | 0.0 | 0.0 | 0.0 | 19.9 | 5.1 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 6.5 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 13.1 | 0.0 | 0.0 | 0.0 | 19.9 | 5.1 |
| Serranidae | | | 0.0 | 6.5 | 2.8 | 14.2 | 2.8 | 5.0 | 1.9 | 1.3 | 0.6 | 4.8 | 9.8 | 10.2 | 60.0 | 15.4 |
| | Epinephelus coioides | Hamour | 0.0 | 6.5 | 2.8 | 14.2 | 2.8 | 5.0 | 1.9 | 1.3 | 0.6 | 4.8 | 9.8 | 10.2 | 60.0 | 15.4 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.8 | 1.5 | 0.7 | 3.5 | 0.9 |
| | Acanthopagrus bifasciatus | Faskar | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| | Argyrops spinifer | Kofar | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 1.5 | 0.7 | 3.0 | 0.8 |
| Sphyraenidae | | | 0.0 | 2.7 | 2.2 | 0.0 | 1.0 | 0.0 | 0.7 | 0.0 | 0.0 | 2.0 | 1.9 | 6.5 | 17.0 | 4.4 |
| | Sphyraena jello | Jedd | 0.0 | 2.7 | 2.2 | 0.0 | 1.0 | 0.0 | 0.7 | 0.0 | 0.0 | 2.0 | 1.9 | 6.5 | 17.0 | 4.4 |

Table 4 (f) - Total catch (tonnes) and species composition for landings at Al Sadar (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|-----------------------|--------------------------------|-----------------|------------|-------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|-------------|
| <i>Tarad</i> | <i>Halaq</i> | | 0.0 | 90.0 | 41.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 131.9 | 34.0 |
| Ariidae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Belonidae et al.</i> | Haqool | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.0 | 0.3 | 8.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.9 | 2.3 |
| | <i>Atule mate</i> | Durduman | 0.0 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.5 | 0.9 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.1 | 5.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.3 | 1.4 |
| Carcharhinidae | | | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| Labridae | | | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.5 |
| | <i>Choerodon robustus</i> | Jin | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.5 |
| Lethrinidae | | | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.2 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Lethrinus nebulosus</i> | Shaari | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.2 |
| Scombridae | | | 0.0 | 85.2 | 31.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 116.7 | 30.0 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.0 | 85.2 | 31.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 116.7 | 30.0 |
| Serranidae | | | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| | <i>Epinephelus coioides</i> | Hamour | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| Sphyraenidae | | | 0.0 | 1.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.7 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 1.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.7 |
| Total | | | 0.0 | 114.8 | 63.1 | 48.9 | 16.8 | 13.0 | 9.9 | 5.7 | 20.1 | 18.5 | 31.3 | 46.2 | 388.3 | 100.0 |
| (%) | | | 0.0 | 29.6 | 16.2 | 12.6 | 4.3 | 3.3 | 2.6 | 1.5 | 5.2 | 4.8 | 8.1 | 11.9 | 100.0 | |

Table 5 - Total number of trips by boat/gear type in the Emirate of Abu Dhabi.

| Landing site | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|--------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|---------------|--------------|
| Al Bateen | 0 | 0 | 1,312 | 640 | 488 | 146 | 883 | 1,293 | 4,761 | 14.2 |
| Al Sadar | 15 | 0 | 0 | 0 | 3,267 | 0 | 457 | 0 | 3,739 | 11.1 |
| Delma Island | 2,375 | 0 | 0 | 0 | 3,188 | 0 | 0 | 0 | 5,563 | 16.5 |
| Free Port | 3,437 | 0 | 0 | 0 | 2,825 | 0 | 0 | 0 | 6,262 | 18.6 |
| Marfa | 0 | 0 | 0 | 0 | 7,203 | 0 | 0 | 0 | 7,203 | 21.4 |
| Mina Zayed | 69 | 0 | 0 | 10 | 2,689 | 0 | 3 | 0 | 2,771 | 8.2 |
| Sila | 370 | 0 | 0 | 0 | 2,634 | 0 | 343 | 0 | 3,347 | 9.9 |
| Total | 6,266 | 0 | 1,312 | 650 | 22,293 | 146 | 1,686 | 1,2930 | 33,646 | 100.0 |
| (%) | 18.6 | 0.0 | 3.9 | 1.9 | 66.3 | 0.4 | 5.0 | 3.8 | 100.0 | |

Table 6 - Total number of trips by landing site in the Emirate of Abu Dhabi.

| Landing site | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|---------------|--------------|
| Al Bateen | 1323 | 904 | 488 | 284 | 157 | 194 | 200 | 195 | 199 | 256 | 342 | 218 | 4,761 | 14.2 |
| <i>Tarad Al Defara</i> | 331 | 82 | 287 | 284 | 22 | 0 | 0 | 49 | 0 | 85 | 171 | 0 | 1,312 | 3.9 |
| <i>Tarad Al Hadhra</i> | 0 | 0 | 0 | 0 | 112 | 151 | 144 | 107 | 125 | 0 | 0 | 0 | 640 | 1.9 |
| <i>Tarad Halaq</i> | 661 | 164 | 57 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 883 | 2.6 |
| <i>Tarad Hadaq</i> | 0 | 0 | 0 | 0 | 22 | 0 | 11 | 0 | 50 | 171 | 171 | 62 | 488 | 1.4 |
| <i>Tarad Leikh</i> | 331 | 657 | 144 | 0 | 0 | 43 | 0 | 0 | 25 | 0 | 0 | 94 | 1,293 | 3.8 |
| <i>Tarad Al Sakkar</i> | 0 | 0 | 0 | 0 | 0 | 0 | 44 | 39 | 0 | 0 | 0 | 62 | 146 | 0.4 |
| Al Sadar | 676 | 390 | 409 | 327 | 214 | 137 | 146 | 118 | 267 | 220 | 426 | 409 | 3,739 | 11.1 |
| <i>Lansh Gargoor</i> | 3 | 6 | 3 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 15 | 0.0 |
| <i>Tarad Halaq</i> | 183 | 157 | 117 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 457 | 1.4 |
| <i>Tarad Hadaq</i> | 490 | 227 | 289 | 327 | 214 | 137 | 146 | 118 | 264 | 220 | 426 | 409 | 3,267 | 9.7 |
| Delma Island | 698 | 465 | 489 | 420 | 400 | 390 | 329 | 248 | 426 | 647 | 540 | 511 | 5,563 | 16.5 |
| <i>Lansh Gargoor</i> | 270 | 98 | 220 | 270 | 189 | 240 | 221 | 140 | 210 | 203 | 147 | 167 | 2,375 | 7.1 |
| <i>Tarad Hadaq</i> | 428 | 367 | 269 | 150 | 211 | 150 | 108 | 108 | 216 | 444 | 393 | 344 | 3,188 | 9.5 |
| Free Port | 890 | 557 | 663 | 750 | 514 | 353 | 358 | 356 | 507 | 573 | 294 | 447 | 6,262 | 18.6 |
| <i>Lansh Gargoor</i> | 341 | 233 | 418 | 477 | 322 | 253 | 227 | 257 | 225 | 322 | 123 | 239 | 3,437 | 10.2 |
| <i>Tarad Hadaq</i> | 549 | 324 | 245 | 273 | 192 | 100 | 131 | 99 | 282 | 251 | 171 | 208 | 2,825 | 8.4 |
| Marfa | 2062 | 1408 | 761 | 431 | 242 | 329 | 317 | 298 | 272 | 317 | 456 | 310 | 7,203 | 21.4 |
| <i>Tarad Hadaq</i> | 2062 | 1408 | 761 | 431 | 242 | 329 | 317 | 298 | 272 | 317 | 456 | 310 | 7,203 | 21.4 |
| Mina Zayed | 430 | 356 | 238 | 324 | 251 | 96 | 138 | 93 | 201 | 175 | 217 | 252 | 2,771 | 8.2 |
| <i>Lansh Gargoor</i> | 7 | 8 | 0 | 3 | 10 | 13 | 7 | 0 | 6 | 8 | 7 | 0 | 69 | 0.2 |
| <i>Tarad Al Hadhra</i> | 0 | 0 | 0 | 0 | 10 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 10 | 0.0 |
| <i>Tarad Halaq</i> | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0.0 |
| <i>Tarad Hadaq</i> | 420 | 348 | 238 | 321 | 231 | 83 | 131 | 93 | 195 | 167 | 210 | 252 | 2,689 | 8.0 |
| Sila | 868 | 631 | 340 | 228 | 144 | 171 | 178 | 124 | 117 | 143 | 224 | 179 | 3,347 | 9.9 |
| <i>Lansh Gargoor</i> | 9 | 9 | 23 | 51 | 46 | 51 | 58 | 28 | 24 | 25 | 22 | 24 | 370 | 1.1 |
| <i>Tarad Halaq</i> | 187 | 106 | 50 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 343 | 1.0 |
| <i>Tarad Hadaq</i> | 672 | 516 | 267 | 177 | 98 | 120 | 120 | 96 | 93 | 118 | 202 | 155 | 2,634 | 7.8 |
| Total | 6,946 | 4,711 | 3,388 | 2,764 | 1,922 | 1,670 | 1,666 | 1,433 | 1,990 | 2,332 | 2,499 | 2,326 | 33,646 | 100.0 |
| (%) | 20.6 | 14.0 | 10.1 | 8.2 | 5.7 | 5.0 | 5.0 | 4.3 | 5.9 | 6.9 | 7.4 | 6.9 | 100.0 | |

Table 7 (a) - Total number of boatdays for lansh by boat/gear type in the Emirate of Abu Dhabi.

| Landing site | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|--------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|---------------|--------------|
| Al Sadar | 36 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 36 | 0.3 |
| Delma Island | 2,533 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2,533 | 17.7 |
| Free Port | 11,291 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 11,291 | 78.8 |
| Mina Zayed | 69 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 69 | 0.5 |
| Sila | 406 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 406 | 2.8 |
| Total | 14,336 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 14,336 | 100.0 |
| (%) | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | |

Table 7 (b) - Total number of boathours for tarad by boat/gear type in the Emirate of Abu Dhabi.

| Landing site | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|--------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|----------------|--------------|
| Al Bateen | 0 | 0 | 6,349 | 2,690 | 2,588 | 801 | 4,171 | 4,841 | 21,440 | 12.2 |
| Al Sadar | 0 | 0 | 0 | 0 | 21,410 | 0 | 3,449 | 0 | 24,859 | 14.2 |
| Delma Island | 0 | 0 | 0 | 0 | 18,453 | 0 | 0 | 0 | 18,453 | 10.5 |
| Free Port | 0 | 0 | 0 | 0 | 36,643 | 0 | 0 | 0 | 36,643 | 20.9 |
| Marfa | 0 | 0 | 0 | 0 | 35,569 | 0 | 0 | 0 | 35,569 | 20.3 |
| Mina Zayed | 0 | 0 | 0 | 90 | 15,964 | 0 | 18 | 0 | 16,072 | 9.2 |
| Sila | 0 | 0 | 0 | 0 | 19,553 | 0 | 2,521 | 0 | 22,074 | 12.6 |
| Total | 0 | 0 | 6,349 | 2,780 | 150,180 | 801 | 10,159 | 4,841 | 175,109 | 100.0 |
| (%) | 0.0 | 0.0 | 3.6 | 1.6 | 85.8 | 0.5 | 5.8 | 2.8 | 100.0 | |

Table 8 (a) - Total number of boatdays for lansh by landing site in the Emirate of Abu Dhabi.

| Landing site | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|----------------------|--------------|------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|------------|------------|---------------|--------------|
| Al Sadar | 3 | 6 | 21 | 0 | 0 | 0 | 0 | 0 | 6 | 0 | 0 | 0 | 36 | 0.3 |
| <i>Lansh Gargoor</i> | 3 | 6 | 21 | 0 | 0 | 0 | 0 | 0 | 6 | 0 | 0 | 0 | 36 | 0.3 |
| Delma Island | 270 | 98 | 220 | 297 | 208 | 264 | 243 | 140 | 210 | 223 | 176 | 184 | 2,533 | 17.7 |
| <i>Lansh Gargoor</i> | 270 | 98 | 220 | 297 | 208 | 264 | 243 | 140 | 210 | 223 | 176 | 184 | 2,533 | 17.7 |
| Free Port | 1,125 | 792 | 1,254 | 1,479 | 998 | 936 | 795 | 874 | 833 | 1,095 | 394 | 717 | 11,291 | 78.8 |
| <i>Lansh Gargoor</i> | 1125 | 792 | 1254 | 1479 | 998 | 936 | 795 | 874 | 833 | 1095 | 394 | 717 | 11,291 | 78.8 |
| Mina Zayed | 7 | 8 | 0 | 3 | 10 | 13 | 7 | 0 | 6 | 8 | 7 | 0 | 69 | 0.5 |
| <i>Lansh Gargoor</i> | 7 | 8 | 0 | 3 | 10 | 13 | 7 | 0 | 6 | 8 | 7 | 0 | 69 | 0.5 |
| Sila | 9 | 14 | 23 | 56 | 46 | 56 | 58 | 28 | 24 | 28 | 22 | 43 | 406 | 2.8 |
| <i>Lansh Gargoor</i> | 9 | 14 | 23 | 56 | 46 | 56 | 58 | 28 | 24 | 28 | 22 | 43 | 406 | 2.8 |
| Total | 1,414 | 918 | 1,518 | 1,835 | 1,262 | 1,269 | 1,103 | 1,042 | 1,079 | 1,354 | 599 | 944 | 14,336 | 100.0 |
| (%) | 9.9 | 6.4 | 10.6 | 0.0 | 8.8 | 8.9 | 7.7 | 7.3 | 7.5 | 9.4 | 4.2 | 6.6 | 100.0 | |

Table 8 (b) - Total number of boathours for tarad by landing site in the Emirate of Abu Dhabi.

| Landing site | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Total | (%) |
|------------------------|---------------|---------------|---------------|---------------|--------------|--------------|--------------|--------------|--------------|---------------|---------------|---------------|----------------|--------------|
| Al Bateen | 4,630 | 4,025 | 2,498 | 1,420 | 650 | 583 | 833 | 1,182 | 1,046 | 1,367 | 2,051 | 1,154 | 21,440 | 12.2 |
| <i>Tarad Al Defara</i> | 992 | 329 | 1579 | 1420 | 112 | 0 | 0 | 293 | 0 | 598 | 1026 | 0 | 6,349 | 3.6 |
| <i>Tarad Al Hadhra</i> | 0 | 0 | 0 | 0 | 426 | 410 | 567 | 664 | 623 | 0 | 0 | 0 | 2,690 | 1.5 |
| <i>Tarad Halaq</i> | 2976 | 822 | 373 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4,171 | 2.4 |
| <i>Tarad Hadaq</i> | 0 | 0 | 0 | 0 | 112 | 0 | 33 | 0 | 274 | 769 | 1026 | 374 | 2,588 | 1.5 |
| <i>Tarad Leikh</i> | 661 | 2875 | 546 | 0 | 0 | 173 | 0 | 0 | 149 | 0 | 0 | 437 | 4,841 | 2.8 |
| <i>Tarad Al Sakkar</i> | 0 | 0 | 0 | 0 | 0 | 0 | 233 | 225 | 0 | 0 | 0 | 343 | 801 | 0.5 |
| Al Sadar | 4,692 | 3,027 | 2,460 | 2,027 | 1,220 | 1,014 | 1,022 | 602 | 1,610 | 1,430 | 2,769 | 2,986 | 24,859 | 14.2 |
| <i>Tarad Halaq</i> | 1409 | 1256 | 784 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3,449 | 2.0 |
| <i>Tarad Hadaq</i> | 3283 | 1771 | 1676 | 2027 | 1220 | 1014 | 1022 | 602 | 1610 | 1430 | 2769 | 2986 | 21,410 | 12.2 |
| Delma Island | 3,338 | 1,945 | 1,614 | 1,035 | 1,055 | 600 | 853 | 508 | 950 | 2,309 | 2,044 | 2,202 | 18,453 | 10.5 |
| <i>Tarad Hadaq</i> | 3338 | 1945 | 1614 | 1035 | 1055 | 600 | 853 | 508 | 950 | 2309 | 2044 | 2202 | 18,453 | 10.5 |
| Free Port | 6,808 | 3,856 | 4,288 | 3,522 | 2,342 | 1,150 | 1,729 | 1,158 | 2,876 | 3,916 | 2,565 | 2,434 | 36,643 | 20.9 |
| <i>Tarad Hadaq</i> | 6808 | 3856 | 4288 | 3522 | 2342 | 1150 | 1729 | 1158 | 2876 | 3916 | 2565 | 2434 | 36,643 | 20.9 |
| Marfa | 9,719 | 7,042 | 3,424 | 2,074 | 954 | 1,752 | 2,185 | 1,454 | 1,617 | 1,227 | 2,279 | 1,843 | 35,569 | 20.3 |
| <i>Tarad Hadaq</i> | 9719 | 7042 | 3424 | 2074 | 954 | 1752 | 2185 | 1454 | 1617 | 1227 | 2279 | 1843 | 35,569 | 20.3 |
| Mina Zayed | 2,958 | 2,366 | 1,499 | 2,536 | 1,037 | 315 | 406 | 623 | 839 | 1,035 | 1,197 | 1,260 | 16,072 | 9.2 |
| <i>Tarad Al Hadhra</i> | 0 | 0 | 0 | 0 | 90 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 90 | 0.1 |
| <i>Tarad Halaq</i> | 18 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 18 | 0.0 |
| <i>Tarad Hadaq</i> | 2940 | 2366 | 1499 | 2536 | 947 | 315 | 406 | 623 | 839 | 1035 | 1197 | 1260 | 15,964 | 9.1 |
| Sila | 7,955 | 3,608 | 2,658 | 903 | 627 | 576 | 948 | 528 | 474 | 956 | 1,353 | 1,488 | 22,074 | 12.6 |
| <i>Tarad Halaq</i> | 1571 | 615 | 335 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2,521 | 1.4 |
| <i>Tarad Hadaq</i> | 6384 | 2993 | 2323 | 903 | 627 | 576 | 948 | 528 | 474 | 956 | 1353 | 1488 | 19,553 | 11.2 |
| Total | 40,100 | 25,868 | 18,441 | 13,517 | 7,885 | 5,990 | 7,977 | 6,055 | 9,413 | 12,240 | 14,258 | 13,365 | 175,109 | 100.0 |
| (%) | 22.9 | 14.8 | 10.5 | 0.0 | 4.5 | 3.4 | 4.6 | 3.5 | 5.4 | 7.0 | 8.1 | 7.6 | 100.0 | |

Table 9 (a) - CPUE in kg/boatday for lansh landings at Sila.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-------------------|----------------------------------|-----------------|------------|------------|--------------|--------------|------------|-------------|------------|--------------|------------|-------------|------------|-------------|--------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 0.0 | 0.0 | 576.0 | 228.0 | 0.0 | 107.0 | 0.0 | 147.5 | 0.0 | 109.0 | 0.0 | 84.0 | 208.6 | 100.0 |
| Carangidae | | | 0.0 | 0.0 | 8.0 | 0.0 | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 1.3 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.3 |
| | <i>Carangoides malabaricus</i> | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.3 |
| | <i>Caranx ignobilis</i> | Gufdar | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.6 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| Haemulidae | | | 0.0 | 0.0 | 420.0 | 199.5 | 0.0 | 93.5 | 0.0 | 145.5 | 0.0 | 92.0 | 0.0 | 84.0 | 172.4 | 82.7 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 420.0 | 199.5 | 0.0 | 93.5 | 0.0 | 145.5 | 0.0 | 83.5 | 0.0 | 84.0 | 171.0 | 82.0 |
| | <i>Plectorhinchus sordidus</i> | Yanam | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.5 | 0.0 | 0.0 | 1.4 | 0.7 |
| Serranidae | | | 0.0 | 0.0 | 146.0 | 28.5 | 0.0 | 5.5 | 0.0 | 2.0 | 0.0 | 17.0 | 0.0 | 0.0 | 33.2 | 15.9 |
| | <i>Epinephelus coioides</i> | Hamour | 0.0 | 0.0 | 146.0 | 28.5 | 0.0 | 5.5 | 0.0 | 2.0 | 0.0 | 17.0 | 0.0 | 0.0 | 33.2 | 15.9 |
| Sparidae | | | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.2 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.2 |
| Total | | | 0.0 | 0.0 | 576.0 | 228.0 | 0.0 | 107.0 | 0.0 | 147.5 | 0.0 | 109.0 | 0.0 | 84.0 | 208.6 | 100.0 |
| (%) | | | 0.0 | 0.0 | 46.0 | 18.2 | 0.0 | 8.5 | 0.0 | 11.8 | 0.0 | 8.7 | 0.0 | 6.7 | 100.0 | |

Table 9 (b) - CPUE in kg/boatday for lansh landings at Delma Island.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|----------------------|---------------------------------|-----------------|-------------|-------------|-------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 169.7 | 209.0 | 139.1 | 405.3 | 397.2 | 412.9 | 322.7 | 257.8 | 228.8 | 230.3 | 313.0 | 371.0 | 288.1 | 100.0 |
| Ariidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.2 | 0.1 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.2 | 0.1 |
| Carangidae | | | 16.0 | 21.7 | 4.8 | 12.7 | 45.3 | 76.5 | 27.9 | 19.4 | 10.5 | 15.5 | 17.9 | 9.8 | 23.2 | 8.0 |
| | <i>Alepes vari</i> | Gamlouh | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.5 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Atule mate</i> | Durduman | 0.0 | 0.0 | 0.0 | 3.4 | 23.3 | 29.1 | 3.5 | 2.1 | 0.6 | 3.8 | 0.7 | 0.3 | 5.6 | 1.9 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 1.5 | 0.0 | 0.0 | 0.0 | 1.4 | 4.6 | 1.7 | 1.1 | 0.3 | 1.1 | 0.4 | 0.3 | 1.0 | 0.4 |
| | <i>Carangoides malabaricus</i> | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 |
| | <i>Caranx heberi</i> | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Caranx ignobilis</i> | Gufdar | 1.6 | 4.3 | 0.4 | 1.3 | 12.9 | 18.8 | 6.7 | 2.8 | 3.7 | 2.1 | 2.7 | 1.6 | 4.9 | 1.7 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 12.6 | 17.3 | 4.3 | 7.9 | 4.6 | 17.6 | 14.9 | 13.0 | 5.8 | 8.4 | 14.1 | 7.4 | 10.7 | 3.7 |
| | <i>Scomberoides</i> | Dhil'e | 0.2 | 0.0 | 0.0 | 0.1 | 1.7 | 0.8 | 0.3 | 0.0 | 0.1 | 0.2 | 0.0 | 0.1 | 0.3 | 0.1 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 5.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.2 |
| | <i>Seriolina nigrofasciata</i> | Hamam | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ephippidae | | | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 4.3 | 13.0 | 3.2 | 1.4 | 0.1 | 2.1 | 0.7 |
| | <i>Ephippidae et al.</i> | Imad | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 4.3 | 13.0 | 3.2 | 1.4 | 0.1 | 2.1 | 0.7 |
| Haemulidae | | | 13.8 | 0.0 | 0.0 | 220.9 | 156.2 | 150.9 | 141.7 | 131.0 | 111.9 | 106.7 | 152.5 | 187.3 | 114.4 | 39.7 |
| | <i>Diagramma pictum</i> | Farsh | 13.8 | 0.0 | 0.0 | 220.9 | 156.0 | 150.7 | 141.6 | 130.8 | 111.9 | 106.7 | 152.5 | 186.8 | 114.3 | 39.7 |
| | <i>Plectorhinchus gaterinus</i> | Hilali | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Plectorhinchus sordidus</i> | Yanam | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.5 | 0.1 | 0.0 |
| Labridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Choerodon robustus</i> | Jin | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Lethrinidae | | | 40.7 | 13.7 | 38.3 | 65.8 | 116.7 | 98.6 | 73.1 | 33.4 | 19.1 | 23.8 | 47.3 | 81.9 | 54.4 | 18.9 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.0 | 0.0 | 0.0 | 3.0 | 2.2 | 1.2 | 1.0 | 0.0 | 0.0 | 0.1 | 0.2 | 0.8 | 0.7 | 0.2 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 8.3 | 9.6 | 7.1 | 5.5 | 3.4 | 1.6 | 2.3 | 3.7 | 10.6 | 4.3 | 1.5 |
| | <i>Lethrinus nebulosus</i> | Shaari | 40.7 | 13.7 | 38.3 | 54.5 | 105.0 | 90.2 | 66.7 | 30.0 | 17.5 | 21.5 | 43.4 | 70.5 | 49.3 | 17.1 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.1 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lutjanus malabaricus</i> | Hamra | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lutjanus russelli</i> | Umm Dhrais | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Pomacanthidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Pomacanthus maculosus</i> | Anfooz | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

Table 9 (b) - CPUE in kg/boatday for lansh landings at Delma Island (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|----------------------------------|-------------|-------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 169.7 | 209.0 | 139.1 | 405.3 | 397.2 | 412.9 | 322.7 | 257.8 | 228.8 | 230.3 | 313.0 | 371.0 | 288.1 | 100.0 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 | 0.0 | 0.1 | 0.0 | 0.8 | 0.6 | 1.9 | 0.3 | 0.1 |
| | <i>Portunus pelagicus</i> | Qabqoob | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 | 0.0 | 0.1 | 0.0 | 0.8 | 0.6 | 1.9 | 0.3 | 0.1 |
| Rachycentridae | | | 0.5 | 0.0 | 0.0 | 0.3 | 1.3 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.3 | 0.1 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.5 | 0.0 | 0.0 | 0.3 | 1.3 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.3 | 0.1 |
| Scombridae | | | 0.5 | 0.0 | 0.0 | 0.1 | 2.1 | 0.6 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 |
| | <i>Rastrelliger kanagurta</i> | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 0.6 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.5 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Serranidae | | | 88.1 | 169.0 | 91.9 | 98.9 | 69.9 | 80.0 | 74.7 | 69.1 | 73.3 | 79.9 | 92.4 | 83.5 | 89.2 | 31.0 |
| | <i>Epinephelus coioides</i> | Hamour | 88.1 | 169.0 | 91.9 | 98.9 | 69.9 | 80.0 | 74.7 | 69.1 | 73.3 | 79.9 | 92.4 | 83.5 | 89.2 | 31.0 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Siganus canaliculatus</i> | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sparidae | | | 8.8 | 4.7 | 4.1 | 6.4 | 3.8 | 4.2 | 1.5 | 0.4 | 0.7 | 0.3 | 0.8 | 4.9 | 3.4 | 1.2 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 8.8 | 4.7 | 4.1 | 6.2 | 3.5 | 3.9 | 1.4 | 0.4 | 0.7 | 0.3 | 0.8 | 4.8 | 3.3 | 1.1 |
| | <i>Acanthopagrus latus</i> | Shaam | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Argyrops spinifer</i> | Kofar | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sphyraenidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Stromateidae | | | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Pampus argenteus</i> | Zubaidi | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Teraponidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Terapon jarbua</i> | Yelly | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tetraodontidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 |
| | <i>Arothron stellatus</i> | Faqel | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 |
| Total | | | 169.7 | 209.0 | 139.1 | 405.3 | 397.2 | 412.9 | 322.7 | 257.8 | 228.8 | 230.3 | 313.0 | 371.0 | 288.1 | 100.0 |
| (%) | | | 4.9 | 6.0 | 4.0 | 11.7 | 11.5 | 11.9 | 9.3 | 7.5 | 6.6 | 6.7 | 9.1 | 10.7 | 100.0 | |

Table 9 (c) - CPUE in kg/boatday for lansh landings at Free Port.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|----------------------------------|-----------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 220.5 | 180.2 | 219.4 | 251.7 | 259.4 | 189.4 | 169.4 | 174.3 | 122.3 | 145.0 | 187.3 | 201.6 | 193.4 | 100.0 |
| Ariidae | | | 0.1 | 1.3 | 0.2 | 0.5 | 1.9 | 1.7 | 0.6 | 0.7 | 0.1 | 0.2 | 2.9 | 2.2 | 1.0 | 0.5 |
| | <i>Arius thalassinus</i> | Khan | 0.1 | 1.3 | 0.2 | 0.5 | 1.9 | 1.7 | 0.6 | 0.7 | 0.1 | 0.2 | 2.9 | 2.2 | 1.0 | 0.5 |
| Belonidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Belonidae et al.</i> | Haqool | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 23.4 | 8.8 | 10.7 | 16.0 | 75.4 | 50.4 | 24.9 | 30.5 | 33.3 | 34.3 | 52.5 | 37.4 | 33.1 | 17.1 |
| | <i>Alepes vari</i> | Gamlouh | 0.1 | 0.0 | 0.0 | 0.1 | 0.2 | 0.2 | 0.5 | 0.1 | 0.1 | 0.3 | 0.3 | 0.1 | 0.1 | 0.1 |
| | <i>Atule mate</i> | Durduman | 2.0 | 0.7 | 0.5 | 3.5 | 29.4 | 17.5 | 5.9 | 10.8 | 13.0 | 18.7 | 15.6 | 4.9 | 10.2 | 5.3 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 3.5 | 0.6 | 1.6 | 1.7 | 2.6 | 5.2 | 4.1 | 2.5 | 7.1 | 3.2 | 6.0 | 5.9 | 3.7 | 1.9 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 1.5 | 1.0 | 1.8 | 7.6 | 2.3 | 1.4 | 1.4 | 0.4 | 3.0 | 10.4 | 1.8 | 2.7 | 1.4 |
| | <i>Carangoides malabaricus</i> | Jesh Sal | 1.8 | 0.7 | 0.0 | 1.3 | 2.4 | 0.1 | 0.1 | 0.2 | 0.2 | 0.0 | 1.3 | 1.4 | 0.8 | 0.4 |
| | <i>Caranx heberi</i> | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.6 | 1.1 | 0.3 | 0.2 | 0.1 | 0.8 | 0.8 | 0.4 | 0.2 |
| | <i>Caranx ignobilis</i> | Gufdar | 0.3 | 0.0 | 0.6 | 0.3 | 1.9 | 4.1 | 1.0 | 5.4 | 0.7 | 0.7 | 5.0 | 2.4 | 1.9 | 1.0 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 14.3 | 4.1 | 5.8 | 5.9 | 21.2 | 13.3 | 9.6 | 7.8 | 11.0 | 7.6 | 12.0 | 16.1 | 10.7 | 5.5 |
| | <i>Megalaspis cordyla</i> | Diyayuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 |
| | <i>Parastromateus niger</i> | Halwayoh | 0.0 | 0.2 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.1 | 0.1 | 0.0 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.0 | 0.1 | 0.3 | 4.6 | 1.7 | 0.0 | 0.2 | 0.0 | 0.5 | 0.6 | 0.5 | 0.7 | 0.4 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 1.6 | 0.6 | 0.6 | 0.7 | 0.1 | 0.0 | 0.7 | 0.4 | 0.2 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.6 | 0.5 | 0.9 | 4.7 | 2.8 | 0.6 | 1.2 | 0.0 | 0.1 | 0.0 | 0.9 | 1.0 | 0.5 |
| | <i>Seriolina nigrofasciata</i> | Hamam | 1.4 | 0.5 | 0.2 | 0.1 | 0.2 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.4 | 1.2 | 0.3 | 0.2 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.3 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.0 | 0.0 | 0.3 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Chanidae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Chanos chanos</i> | Aifah | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Clupeidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Sardinella longiceps</i> | Oumah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Ephippidae | | | 0.0 | 0.1 | 0.1 | 0.0 | 0.1 | 1.6 | 5.0 | 3.4 | 4.5 | 3.4 | 0.9 | 0.9 | 1.7 | 0.9 |
| | <i>Ephippidae et al.</i> | Imad | 0.0 | 0.1 | 0.1 | 0.0 | 0.1 | 1.6 | 5.0 | 3.4 | 4.5 | 3.4 | 0.9 | 0.9 | 1.7 | 0.9 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Gerres longirostris</i> | Badah | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Haemulidae | | | 31.3 | 32.5 | 38.7 | 29.4 | 30.4 | 20.7 | 22.3 | 26.9 | 15.4 | 20.6 | 30.5 | 37.4 | 28.0 | 14.5 |
| | <i>Diagramma pictum</i> | Farsh | 17.9 | 19.0 | 25.7 | 25.9 | 27.2 | 19.2 | 20.5 | 24.8 | 13.7 | 17.9 | 25.1 | 29.4 | 22.2 | 11.5 |
| | <i>Plectorhinchus gaterinus</i> | Hilali | 5.4 | 1.9 | 2.7 | 0.6 | 1.8 | 1.1 | 1.0 | 1.6 | 1.1 | 1.8 | 3.6 | 4.7 | 2.3 | 1.2 |
| | <i>Plectorhinchus sordidus</i> | Yanam | 8.0 | 11.6 | 10.1 | 2.7 | 1.4 | 0.4 | 0.8 | 0.5 | 0.6 | 0.9 | 1.8 | 3.2 | 3.5 | 1.8 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.2 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

Table 9 (c) - CPUE in kg/boatday for lansh landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|--------------------------|-------------------------|-----------------|-------------|-------------|-------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 220.5 | 180.2 | 219.4 | 251.7 | 259.4 | 189.4 | 169.4 | 174.3 | 122.3 | 145.0 | 187.3 | 201.6 | 193.4 | 100.0 |
| Labridae | | | 0.0 | 0.4 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 |
| | Choerodon robustus | Jin | 0.0 | 0.4 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 |
| Lethrinidae | | | 25.3 | 30.4 | 48.8 | 105.0 | 85.8 | 57.4 | 57.4 | 52.1 | 21.9 | 28.9 | 45.2 | 33.7 | 49.3 | 25.5 |
| | Lethrinus borbonicus | Yemah | 3.4 | 3.8 | 4.6 | 10.0 | 2.2 | 6.0 | 4.9 | 3.7 | 2.7 | 1.8 | 2.1 | 3.0 | 4.0 | 2.1 |
| | Lethrinus lentjan | Shaari Eshkheli | 4.0 | 3.9 | 6.8 | 14.2 | 11.8 | 11.7 | 18.8 | 16.1 | 4.3 | 9.8 | 17.2 | 11.6 | 10.8 | 5.6 |
| | Lethrinus microdon | Souli | 1.1 | 2.7 | 1.3 | 1.0 | 1.6 | 1.9 | 2.5 | 2.8 | 1.4 | 1.5 | 1.8 | 1.2 | 1.7 | 0.9 |
| | Lethrinus nebulosus | Shaari | 16.8 | 20.2 | 36.1 | 79.8 | 70.3 | 37.8 | 31.3 | 29.5 | 13.5 | 15.8 | 24.2 | 17.8 | 32.7 | 16.9 |
| Lutjanidae | | | 0.3 | 0.3 | 0.8 | 0.8 | 1.4 | 1.3 | 0.8 | 0.7 | 1.0 | 0.8 | 0.7 | 0.9 | 0.8 | 0.4 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.0 | 0.0 | 0.5 | 0.3 | 0.8 | 0.1 | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 0.2 | 0.1 |
| | Lutjanus fulviflamma | Aqalah | 0.2 | 0.1 | 0.8 | 0.2 | 1.0 | 0.3 | 0.4 | 0.6 | 0.8 | 0.7 | 0.2 | 0.0 | 0.4 | 0.2 |
| | Lutjanus malabaricus | Hamra | 0.1 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 0.1 | 0.2 | 0.1 | 0.3 | 0.6 | 0.1 | 0.1 |
| | Lutjanus russelli | Umm Dhrais | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 | 0.0 | 0.0 |
| Mixed | | | 0.3 | 0.7 | 0.5 | 0.8 | 0.9 | 0.6 | 0.8 | 0.5 | 0.4 | 0.4 | 0.4 | 1.0 | 0.6 | 0.3 |
| | Mixed species | Mekhlot | 0.3 | 0.7 | 0.5 | 0.8 | 0.9 | 0.6 | 0.8 | 0.5 | 0.4 | 0.4 | 0.4 | 1.0 | 0.6 | 0.3 |
| Myliobatidae etc. | | | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Myliobatidea et al. | Lugmah | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Nemipteridae | | | 2.9 | 5.7 | 4.9 | 3.0 | 0.9 | 2.0 | 0.7 | 0.8 | 1.6 | 0.9 | 1.9 | 4.3 | 2.5 | 1.3 |
| | Scolopsis taeniatus | Ebzimi | 2.9 | 5.7 | 4.9 | 3.0 | 0.9 | 2.0 | 0.7 | 0.8 | 1.6 | 0.9 | 1.9 | 4.3 | 2.5 | 1.3 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Pomacanthidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Pomacanthus maculosus | Anfooz | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Portunidae | | | 0.8 | 0.3 | 0.1 | 0.6 | 0.4 | 0.5 | 1.0 | 1.5 | 2.2 | 5.2 | 6.9 | 6.8 | 2.2 | 1.1 |
| | Portunus pelagicus | Qabqoob | 0.8 | 0.3 | 0.1 | 0.6 | 0.4 | 0.5 | 1.0 | 1.5 | 2.2 | 5.2 | 6.9 | 6.8 | 2.2 | 1.1 |
| Rachycentridae | | | 1.8 | 3.4 | 2.4 | 1.5 | 4.6 | 1.6 | 0.7 | 1.5 | 0.1 | 2.5 | 3.3 | 3.7 | 2.3 | 1.2 |
| | Rachycentron canadum | Sikkil | 1.8 | 3.4 | 2.4 | 1.5 | 4.6 | 1.6 | 0.7 | 1.5 | 0.1 | 2.5 | 3.3 | 3.7 | 2.3 | 1.2 |
| Scaridae | | | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 1.1 | 0.3 | 0.3 | 0.7 | 0.0 | 0.3 | 0.9 | 0.3 | 0.2 |
| | Scarus ghobban | Gain | 0.0 | 0.1 | 0.1 | 0.0 | 0.2 | 1.1 | 0.3 | 0.3 | 0.7 | 0.0 | 0.3 | 0.9 | 0.3 | 0.2 |
| Scombridae | | | 0.0 | 1.7 | 6.2 | 0.8 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.9 | 0.2 | 0.0 | 0.8 | 0.4 |
| | Euthynnus affinis | Sadah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 |
| | Scomberomorus commerson | Kanaad | 0.0 | 1.7 | 6.2 | 0.7 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.9 | 0.0 | 0.0 | 0.8 | 0.4 |

Table 9 (c) - CPUE in kg/boatday for lansh landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|---------------------|---------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 220.5 | 180.2 | 219.4 | 251.7 | 259.4 | 189.4 | 169.4 | 174.3 | 122.3 | 145.0 | 187.3 | 201.6 | 193.4 | 100.0 |
| Serranidae | | | 93.4 | 68.1 | 88.1 | 82.6 | 47.4 | 45.8 | 50.2 | 50.2 | 36.6 | 42.3 | 36.3 | 59.2 | 58.4 | 30.2 |
| | Cephalopholis hemistiktos | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Epinephelus areolatus | Semman | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Epinephelus coioides | Hamour | 93.4 | 68.1 | 88.1 | 82.5 | 47.4 | 45.8 | 50.1 | 50.1 | 36.6 | 42.3 | 36.3 | 59.2 | 58.3 | 30.2 |
| Siganidae | | | 0.4 | 0.6 | 0.7 | 2.9 | 1.9 | 0.9 | 1.4 | 1.7 | 1.8 | 1.8 | 1.5 | 0.3 | 1.3 | 0.7 |
| | Siganus canaliculatus | Safi Arabi | 0.4 | 0.6 | 0.7 | 2.9 | 1.9 | 0.9 | 1.4 | 1.7 | 1.8 | 1.8 | 1.5 | 0.3 | 1.3 | 0.7 |
| | Siganus javus | Safi Sneifi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sparidae | | | 39.8 | 25.7 | 17.2 | 7.3 | 7.5 | 3.4 | 3.1 | 3.0 | 2.5 | 2.2 | 3.7 | 12.8 | 10.7 | 5.5 |
| | Acanthopagrus bifasciatus | Faskar | 9.2 | 8.4 | 8.5 | 1.6 | 2.9 | 2.4 | 2.8 | 2.8 | 2.4 | 1.9 | 3.1 | 4.5 | 4.2 | 2.2 |
| | Acanthopagrus latus | Shaam | 0.5 | 0.7 | 1.2 | 0.6 | 0.2 | 0.3 | 0.1 | 0.1 | 0.0 | 0.3 | 0.1 | 0.5 | 0.4 | 0.2 |
| | Argyrops spinifer | Kofar | 12.7 | 9.5 | 7.5 | 5.0 | 4.4 | 0.7 | 0.1 | 0.0 | 0.1 | 0.0 | 0.6 | 2.5 | 3.6 | 1.9 |
| | Rhabdosargus sarba | Qabit | 17.4 | 7.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.4 | 2.5 | 1.3 |
| Sphyraenidae | | | 0.0 | 0.1 | 0.0 | 0.1 | 0.1 | 0.3 | 0.0 | 0.3 | 0.1 | 0.6 | 0.0 | 0.0 | 0.1 | 0.1 |
| | Sphyraena jello | Jedd | 0.0 | 0.1 | 0.0 | 0.1 | 0.1 | 0.3 | 0.0 | 0.3 | 0.1 | 0.6 | 0.0 | 0.0 | 0.1 | 0.1 |
| Stromateidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Pampus argenteus | Zubaidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total | | | 220.5 | 180.2 | 219.4 | 251.7 | 259.4 | 189.4 | 169.4 | 174.3 | 122.3 | 145.0 | 187.3 | 201.6 | 193.4 | 100.0 |
| (%) | | | 9.5 | 7.8 | 9.5 | 10.8 | 11.2 | 8.2 | 7.3 | 7.5 | 5.3 | 6.2 | 8.1 | 8.7 | 100.0 | |

Table 9 (d) - CPUE in kg/boatday for lanch landings at Al Sadar.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|----------------------------------|-----------------|------------|------------|------------|------------|------------|-------------|------------|------------|------------|------------|------------|------------|-------------|-------------|
| <i>Lansh</i> | <i>Gargoor</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 262.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 262.3 | 100.0 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 | 3.4 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.7 | 2.5 |
| | <i>Carangoides malabaricus</i> | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.3 | 0.9 |
| Ephippidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.6 |
| | <i>Ephippidae et al.</i> | Imad | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.6 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 36.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 36.7 | 14.0 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 36.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 36.7 | 14.0 |
| Lethrinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 93.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 93.3 | 35.6 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 1.9 |
| | <i>Lethrinus lentjan</i> | Shaari Eshkheli | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 40.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 40.0 | 15.2 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 1.9 |
| | <i>Lethrinus nebulosus</i> | Shaari | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 43.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 43.3 | 16.5 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.3 | 1.3 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.3 | 1.3 |
| Nemipteridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 1.9 |
| | <i>Scolopsis taeniatus</i> | Ebzimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 1.9 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 13.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 13.3 | 5.1 |
| | <i>Portunus pelagicus</i> | Qabqoob | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 13.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 13.3 | 5.1 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.3 | 3.2 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.3 | 3.2 |
| Serranidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 90.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 90.0 | 34.3 |
| | <i>Epinephelus coioides</i> | Hamour | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 90.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 90.0 | 34.3 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.6 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.6 |
| Total | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 262.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 262.3 | 100.0 |
| (%) | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | |

Table 10 (a) - CPUE in kg/boathour for tarad landings at Sila.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|-------------------------|-------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 6.2 | 8.9 | 5.5 | 10.0 | 9.2 | 19.1 | 5.1 | 11.9 | 9.7 | 9.4 | 13.6 | 7.4 | 9.7 | 82.6 |
| Carangidae | | | 0.1 | 0.3 | 0.7 | 4.5 | 7.2 | 5.5 | 0.6 | 1.0 | 4.9 | 2.1 | 7.9 | 1.0 | 3.0 | 25.6 |
| | Atule mate | Durduman | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.2 |
| | Carangidae all | Jesh unidentified | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.2 | 0.4 | 2.1 | 5.9 | 3.0 | 0.4 | 0.9 | 4.9 | 1.7 | 7.8 | 0.9 | 2.4 | 20.1 |
| | Carangoides malabaricus | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.1 |
| | Caranx ignobilis | Gufdar | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.4 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.7 |
| | Gnathanodon speciosus | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 |
| | Scomberoides tol | Bassar | 0.0 | 0.1 | 0.3 | 2.3 | 0.8 | 0.1 | 0.3 | 0.0 | 0.0 | 0.1 | 0.1 | 0.0 | 0.3 | 2.9 |
| Carcharhinidae | | | 0.0 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.4 |
| | Carcharhinidae et al. | Jarjoor | 0.0 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.4 |
| Chanidae | | | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| | Chanos chanos | Aifah | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Haemulidae | | | 0.0 | 0.3 | 0.2 | 2.3 | 0.3 | 9.1 | 0.6 | 5.9 | 3.0 | 2.6 | 0.0 | 3.6 | 2.3 | 19.9 |
| | Diagramma pictum | Farsh | 0.0 | 0.3 | 0.2 | 2.3 | 0.3 | 9.1 | 0.6 | 5.9 | 3.0 | 2.6 | 0.0 | 3.6 | 2.3 | 19.9 |
| Lethrinidae | | | 0.0 | 0.9 | 0.3 | 1.7 | 0.6 | 0.3 | 0.9 | 0.0 | 0.0 | 0.4 | 1.3 | 1.3 | 0.6 | 5.5 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| | Lethrinus microdon | Souli | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.1 |
| | Lethrinus nebulosus | Shaari | 0.0 | 0.8 | 0.3 | 1.2 | 0.6 | 0.3 | 0.9 | 0.0 | 0.0 | 0.4 | 1.3 | 1.3 | 0.6 | 5.0 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| Rachycentridae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Rachycentron canadum | Sikkil | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Scombridae | | | 5.7 | 6.4 | 4.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 11.5 |
| | Scomberomorus commerson | Kanaad | 5.7 | 6.4 | 4.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 11.5 |
| Serranidae | | | 0.1 | 0.1 | 0.2 | 1.2 | 0.8 | 4.2 | 2.9 | 4.9 | 1.8 | 4.3 | 4.3 | 1.4 | 2.2 | 18.7 |
| | Epinephelus coioides | Hamour | 0.1 | 0.1 | 0.2 | 1.2 | 0.8 | 4.2 | 2.9 | 4.9 | 1.8 | 4.3 | 4.3 | 1.4 | 2.2 | 18.7 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sphyraenidae | | | 0.1 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| | Sphyraena jello | Jedd | 0.1 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |

Table 10 (a) - CPUE in kg/boathour for tarad landings at Sila (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|--------------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Halaq</i> | | 4.7 | 9.8 | 9.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 17.4 |
| Ariidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.5 | 0.4 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.2 |
| | <i>Atule mate</i> | Durduman | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.2 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 |
| | <i>Caranx ignobilis</i> | Gufdar | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Scomberoides tol</i> | Bassar | 0.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.5 |
| Carcharhinidae | | | 0.0 | 0.5 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.5 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.5 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.5 |
| Chanidae | | | 0.1 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| | <i>Chanos chanos</i> | Aifah | 0.1 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| Haemulidae | | | 0.4 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.6 |
| | <i>Diagramma pictum</i> | Farsh | 0.4 | 1.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.6 |
| | <i>Plectorhinchus sordidus</i> | Yanam | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Lethrinidae | | | 0.1 | 1.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.0 |
| | <i>Lethrinus nebulosus</i> | Shaari | 0.1 | 1.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.0 |
| Rachycentridae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Scombridae | | | 2.7 | 4.5 | 8.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 11.0 |
| | <i>Scomberomorus commerson</i> | Kanaad | 2.7 | 4.5 | 8.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 11.0 |
| Serranidae | | | 0.9 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.5 |
| | <i>Epinephelus coioides</i> | Hamour | 0.9 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 1.5 |
| Sphyraenidae | | | 0.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| | <i>Sphyraena jello</i> | Jedd | 0.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| Total | | | 10.9 | 18.7 | 15.4 | 10.0 | 9.2 | 19.1 | 5.1 | 11.9 | 9.7 | 9.4 | 13.6 | 7.4 | 11.7 | 100.0 |
| (%) | | | 7.8 | 13.3 | 11.0 | 7.2 | 6.5 | 13.6 | 3.6 | 8.5 | 6.9 | 6.7 | 9.7 | 5.3 | 100.0 | |

Table 10 (b) - CPUE in kg/boathour for tarad landings at Delma Island.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|--------------------|----------------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|------------|------------|-------------|-------------|-------------|
| <i>Tarad</i> | <i>Gargoor</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 127.0 | 0.0 | 0.0 | 294.0 | 35.1 | 81.2 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 32.0 | 3.4 | 7.9 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 0.8 | 1.7 |
| | <i>Caranx ignobilis</i> | Gufdar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.4 | 1.0 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 27.0 | 2.3 | 5.2 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 86.0 | 0.0 | 0.0 | 80.0 | 13.8 | 32.0 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 86.0 | 0.0 | 0.0 | 80.0 | 13.8 | 32.0 |
| Lethrinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 85.0 | 7.4 | 17.2 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.4 | 1.0 |
| | <i>Lethrinus nebulosus</i> | Shaari | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 80.0 | 7.0 | 16.2 |
| Serranidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.0 | 0.0 | 0.0 | 95.0 | 10.3 | 23.7 |
| | <i>Epinephelus coioides</i> | Hamour | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.0 | 0.0 | 0.0 | 95.0 | 10.3 | 23.7 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.2 | 0.4 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.2 | 0.4 |

Table 10 (b) - CPUE in kg/boathour for tarad landings at Delma Island (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|----------------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 12.1 | 2.0 | 11.1 | 9.0 | 6.0 | 4.2 | 6.0 | 5.1 | 5.9 | 9.3 | 9.9 | 11.3 | 7.7 | 17.7 |
| Ariidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Belonidae et al.</i> | Haqool | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 1.6 | 0.0 | 1.8 | 5.2 | 3.7 | 0.8 | 2.9 | 1.5 | 3.4 | 1.6 | 0.9 | 0.4 | 2.0 | 4.6 |
| | <i>Atule mate</i> | Durduman | 0.0 | 0.0 | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.8 | 0.0 | 0.0 | 3.6 | 3.3 | 0.4 | 2.7 | 0.9 | 1.5 | 0.5 | 0.7 | 0.4 | 1.2 | 2.8 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Caranx ignobilis</i> | Gufdar | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.5 | 0.0 | 1.8 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.5 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.0 | 0.0 | 0.6 | 0.2 | 0.3 | 0.2 | 0.3 | 1.5 | 1.1 | 0.3 | 0.0 | 0.4 | 0.8 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.2 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Carcharhinidae | | | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Haemulidae | | | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Diagramma pictum</i> | Farsh | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Lethrinidae | | | 4.7 | 2.0 | 5.6 | 2.6 | 0.7 | 0.8 | 0.9 | 0.2 | 0.2 | 0.4 | 2.6 | 8.4 | 2.4 | 5.6 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.3 | 0.1 | 0.1 | 0.1 | 0.2 | 0.6 | 0.1 | 0.3 |
| | <i>Lethrinus nebulosus</i> | Shaari | 4.7 | 2.0 | 5.6 | 2.6 | 0.6 | 0.7 | 0.5 | 0.1 | 0.1 | 0.3 | 2.4 | 7.8 | 2.3 | 5.3 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Rachycentridae | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Scombridae | | | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 | 2.2 | 1.6 | 2.8 | 2.0 | 6.6 | 6.1 | 1.9 | 2.1 | 4.8 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 | 2.2 | 1.6 | 2.8 | 2.0 | 6.6 | 6.1 | 1.9 | 2.1 | 4.8 |
| Serranidae | | | 3.8 | 0.0 | 3.7 | 0.4 | 0.1 | 0.2 | 0.5 | 0.0 | 0.2 | 0.1 | 0.2 | 0.3 | 0.8 | 1.8 |
| | <i>Cephalopholis hemistiktos</i> | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Epinephelus coioides</i> | Hamour | 3.8 | 0.0 | 3.7 | 0.4 | 0.1 | 0.2 | 0.5 | 0.0 | 0.2 | 0.1 | 0.2 | 0.3 | 0.8 | 1.8 |
| Sparidae | | | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.3 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Rhabdosargus sarba</i> | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.2 |
| Sphyraenidae | | | 0.1 | 0.0 | 0.0 | 0.8 | 0.6 | 0.2 | 0.1 | 0.0 | 0.1 | 0.0 | 0.1 | 0.1 | 0.2 | 0.4 |
| | <i>Sphyraena jello</i> | Jedd | 0.1 | 0.0 | 0.0 | 0.8 | 0.6 | 0.2 | 0.1 | 0.0 | 0.1 | 0.0 | 0.1 | 0.1 | 0.2 | 0.4 |

Table 10 (b) - CPUE in kg/boathour for tarad landings at Delma Island (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|--------------------|-----------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Halaq</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 1.1 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 1.0 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.8 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| Lethrinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Serranidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Epinephelus coioides</i> | Hamour | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total | | | 12.1 | 2.0 | 11.1 | 9.0 | 6.0 | 4.2 | 6.0 | 10.6 | 132.9 | 9.3 | 9.9 | 305.3 | 43.2 | 100.0 |
| (%) | | | 2.3 | 0.4 | 2.1 | 1.7 | 1.2 | 0.8 | 1.2 | 2.0 | 25.6 | 1.8 | 1.9 | 58.9 | 100.0 | |

Table 10 (c) - CPUE in kg/boathour for tarad landings at Marfa.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|----------------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 13.2 | 14.0 | 12.6 | 24.1 | 13.8 | 12.8 | 9.7 | 9.5 | 12.5 | 12.9 | 11.3 | 15.5 | 13.5 | 100.0 |
| Ariidae | | | 0.2 | 0.3 | 0.0 | 1.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 1.2 |
| | <i>Arius thalassinus</i> | Khan | 0.2 | 0.3 | 0.0 | 1.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 1.2 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Belonidae et al.</i> | Haqool | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Carangidae | | | 4.8 | 5.1 | 4.6 | 5.9 | 3.7 | 2.5 | 4.8 | 2.9 | 0.3 | 4.2 | 2.5 | 2.7 | 3.7 | 27.3 |
| | <i>Atule mate</i> | Durduman | 0.6 | 1.3 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 1.5 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.1 | 0.0 | 0.2 | 5.8 | 0.5 | 2.1 | 3.4 | 2.7 | 0.1 | 1.1 | 0.0 | 0.4 | 1.4 | 10.1 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 4.2 | 3.8 | 4.4 | 0.0 | 3.2 | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 2.3 | 2.3 | 1.9 | 14.2 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.2 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 1.3 | 0.3 | 0.2 | 0.0 | 0.1 | 0.0 | 0.2 | 1.2 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.4 | 0.6 | 0.8 | 0.0 | 0.0 | 0.0 | 0.3 | 1.9 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.6 | 0.8 | 0.0 | 0.0 | 0.0 | 0.2 | 1.4 |
| | <i>Plectorhinchus gaterinus</i> | Hilali | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| Labridae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.4 |
| | <i>Choerodon robustus</i> | Jin | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.4 |
| Lethrinidae | | | 2.1 | 3.3 | 1.1 | 8.1 | 5.6 | 5.2 | 1.7 | 3.3 | 2.9 | 2.9 | 3.8 | 7.0 | 3.9 | 29.0 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.2 | 0.3 | 0.1 | 0.0 | 0.6 | 0.0 | 0.0 | 0.1 | 0.0 | 0.3 | 0.2 | 0.0 | 0.2 | 1.1 |
| | <i>Lethrinus lentjan</i> | Shaari Eshkheli | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.1 | 0.0 | 0.3 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.4 |
| | <i>Lethrinus nebulosus</i> | Shaari | 1.9 | 3.0 | 1.0 | 8.1 | 4.7 | 5.2 | 1.7 | 3.2 | 2.9 | 2.1 | 3.6 | 6.9 | 3.7 | 27.2 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |

Table 10 (c) - CPUE in kg/boathour for tarad landings at Marfa (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|---------------------|---------------------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 13.2 | 14.0 | 12.6 | 24.1 | 13.8 | 12.8 | 9.7 | 9.5 | 12.5 | 12.9 | 11.3 | 15.5 | 13.5 | 100.0 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.6 |
| | Scarus ghobban | Gain | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.6 |
| Scombridae | | | 2.1 | 0.0 | 3.9 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 8.1 | 0.0 | 0.0 | 0.0 | 1.2 | 8.9 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Scomberomorus commerson | Kanaad | 2.1 | 0.0 | 3.9 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 8.1 | 0.0 | 0.0 | 0.0 | 1.2 | 8.9 |
| Serranidae | | | 2.6 | 3.7 | 1.7 | 7.0 | 2.3 | 5.0 | 1.9 | 2.2 | 0.3 | 3.4 | 3.6 | 3.4 | 3.1 | 22.8 |
| | Epinephelus coioides | Hamour | 2.6 | 3.7 | 1.7 | 7.0 | 2.3 | 5.0 | 1.9 | 2.2 | 0.3 | 3.4 | 3.6 | 3.4 | 3.1 | 22.8 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.5 | 0.6 | 0.2 | 0.1 | 1.0 |
| | Acanthopagrus bifasciatus | Faskar | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Argyrops spinifer | Kofar | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.6 | 0.2 | 0.1 | 0.8 |
| Sphyraenidae | | | 1.4 | 1.5 | 1.3 | 0.0 | 0.8 | 0.0 | 0.7 | 0.0 | 0.0 | 1.4 | 0.7 | 2.2 | 0.8 | 6.2 |
| | Sphyraena jello | Jedd | 1.4 | 1.5 | 1.3 | 0.0 | 0.8 | 0.0 | 0.7 | 0.0 | 0.0 | 1.4 | 0.7 | 2.2 | 0.8 | 6.2 |
| Total | | | 13.2 | 14.0 | 12.6 | 24.1 | 13.8 | 12.8 | 9.7 | 9.5 | 12.5 | 12.9 | 11.3 | 15.5 | 13.5 | 100.0 |
| (%) | | | 8.2 | 8.6 | 7.8 | 14.9 | 8.5 | 7.9 | 6.0 | 5.9 | 7.7 | 8.0 | 7.0 | 9.6 | 100.0 | |

Table 10 (d) - CPUE in kg/boathour for tarad landings at Al Bateen.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|------------------------------|-------------|-------------|-------------|------------|-------------|-------------|------------|------------|------------|------------|-------------|-------------|------------|------------|------------|
| <i>Tarad</i> | <i>Al Defara</i> | | 12.3 | 10.0 | 17.1 | 26.0 | 24.0 | 0.0 | 0.0 | 25.7 | 0.0 | 17.9 | 91.7 | 0.0 | 18.7 | 32.8 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 5.8 | 0.0 | 0.7 | 1.1 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 5.8 | 0.0 | 0.7 | 1.1 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 12.9 | 0.0 | 0.0 | 1.1 | 1.9 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 12.9 | 0.0 | 0.0 | 1.1 | 1.9 |
| Chanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 24.0 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.6 | 4.5 |
| | <i>Chanos chanos</i> | Aifah | 0.0 | 0.0 | 0.0 | 0.0 | 24.0 | 0.0 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.6 | 4.5 |
| Gerreidae | | | 12.3 | 10.0 | 8.4 | 0.0 | 0.0 | 0.0 | 0.0 | 3.2 | 0.0 | 0.0 | 5.8 | 0.0 | 3.3 | 5.8 |
| | <i>Gerres longirostris</i> | Badah | 12.3 | 10.0 | 8.4 | 0.0 | 0.0 | 0.0 | 0.0 | 3.2 | 0.0 | 0.0 | 5.8 | 0.0 | 3.3 | 5.8 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 21.7 | 0.0 | 1.8 | 3.2 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 21.7 | 0.0 | 1.8 | 3.2 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| Mugilidae | | | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 58.3 | 0.0 | 5.3 | 9.3 |
| | <i>Valamugil seheli</i> | Beyah Arabi | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 58.3 | 0.0 | 5.3 | 9.3 |
| Portunidae | | | 0.0 | 0.0 | 7.5 | 26.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 3.2 | 5.6 |
| | <i>Portunus pelagicus</i> | Qabqoob | 0.0 | 0.0 | 7.5 | 26.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 3.2 | 5.6 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| | <i>Siganus canaliculatus</i> | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.8 |
| | <i>Rhabdosargus sarba</i> | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.8 |

Table 10 (d) - CPUE in kg/boathour for tarad landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-------------------|-------------------------|-----------------|------------|------------|------------|------------|-------------|-------------|-------------|-------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Al Hadhra</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 26.8 | 28.4 | 19.4 | 20.6 | 17.4 | 0.0 | 0.0 | 0.0 | 9.4 | 16.4 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Belonidae et al. | Haqool | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Scomberoides tol | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Chanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| | Chanos chanos | Aifah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 1.0 | 4.8 | 0.0 | 0.0 | 0.0 | 0.6 | 1.0 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 1.0 | 4.8 | 0.0 | 0.0 | 0.0 | 0.6 | 1.0 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 23.4 | 24.7 | 15.3 | 11.1 | 3.2 | 0.0 | 0.0 | 0.0 | 6.5 | 11.3 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 23.4 | 24.7 | 15.3 | 11.1 | 3.2 | 0.0 | 0.0 | 0.0 | 6.5 | 11.3 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 1.1 | 0.0 | 2.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 1.0 |
| | Portunus pelagicus | Qabqoob | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 1.1 | 0.0 | 2.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 1.0 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.2 | 0.0 | 0.0 | 0.0 | 0.7 | 1.2 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 8.2 | 0.0 | 0.0 | 0.0 | 0.7 | 1.2 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 2.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.5 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 2.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.5 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |

Table 10 (d) - CPUE in kg/boathour for tarad landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|---------------------|-------------------------|-----------------|------------|------------|------------|------------|-------------|------------|-------------|------------|-------------|-------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 20.0 | 0.0 | 10.0 | 0.0 | 19.1 | 28.9 | 7.0 | 7.9 | 7.7 | 13.5 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.3 | 2.2 |
| | Atule mate | Durduman | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.7 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 1.5 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| | Plectorhinchus sordidus | Yanam | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| Lethrinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.8 | 0.2 | 0.3 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 |
| | Lethrinus nebulosus | Shaari | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.7 | 0.1 | 0.2 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 19.1 | 0.0 | 5.0 | 0.0 | 2.0 | 3.5 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 19.1 | 0.0 | 5.0 | 0.0 | 2.0 | 3.5 |
| Serranidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 1.0 | 1.7 |
| | Epinephelus coioides | Hamour | 0.0 | 0.0 | 0.0 | 0.0 | 11.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 1.0 | 1.7 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.9 | 0.0 | 5.0 | 2.8 | 4.9 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 28.9 | 0.0 | 5.0 | 2.8 | 4.9 |
| Sphyraenidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.3 | 0.2 | 0.3 |
| | Sphyraena jello | Jedd | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.3 | 0.2 | 0.3 |

Table 10 (d) - CPUE in kg/boathour for tarad landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|---------------------|----------------------------------|-------------|------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Halaq</i> | | 6.7 | 37.0 | 15.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.9 | 8.6 |
| Carangidae | | | 0.0 | 8.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 1.2 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.5 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 4.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.7 |
| Scombridae | | | 6.7 | 29.0 | 9.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 6.5 |
| | <i>Scomberomorus commerson</i> | Kanaad | 6.7 | 29.0 | 9.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 6.5 |
| Sphyraenidae | | | 0.0 | 0.0 | 6.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.9 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 0.0 | 6.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.9 |

Table 10 (d) - CPUE in kg/boathour for tarad landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|------------------------|---------------------------|-----------------|-------------|------------|-------------|------------|------------|-------------|------------|------------|------------|------------|------------|-------------|------------|------------|
| <i>Tarad</i> | <i>Leikh</i> | | 17.5 | 21.8 | 13.8 | 0.0 | 0.0 | 35.1 | 0.0 | 0.0 | 12.5 | 0.0 | 0.0 | 37.6 | 11.5 | 20.2 |
| Carangidae | | | 0.0 | 4.1 | 0.0 | 0.0 | 0.0 | 3.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.8 | 1.4 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.4 |
| | Carangoides fulvoguttatus | Jesh Zeria | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.5 |
| | Scomberoides | Dhil'e | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| | Scomberoides tol | Bassar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.1 | 0.2 |
| Gerreidae | | | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 4.2 | 0.0 | 0.0 | 18.4 | 2.1 | 3.7 |
| | Gerres longirostris | Badah | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 4.2 | 0.0 | 0.0 | 18.4 | 2.1 | 3.7 |
| Lethrinidae | | | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| | Lethrinus nebulosus | Shaari | 0.0 | 2.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 18.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.1 | 2.2 | 3.8 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 18.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.1 | 2.2 | 3.8 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 | 0.1 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 | 0.1 |
| Portunidae | | | 17.5 | 8.2 | 13.8 | 0.0 | 0.0 | 3.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.5 | 4.2 | 7.4 |
| | Portunus pelagicus | Qabqoob | 17.5 | 8.2 | 13.8 | 0.0 | 0.0 | 3.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 7.5 | 4.2 | 7.4 |
| Scombridae | | | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 7.5 | 0.0 | 0.0 | 8.3 | 0.0 | 0.0 | 0.0 | 1.4 | 2.4 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 7.5 | 0.0 | 0.0 | 8.3 | 0.0 | 0.0 | 0.0 | 1.4 | 2.4 |
| Serranidae | | | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 | 0.2 |
| | Epinephelus coioides | Hamour | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 | 0.2 |
| Siganidae | | | 0.0 | 3.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.4 | 0.7 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 3.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.4 | 0.7 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 | 0.1 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 | 0.1 |
| Sphyraenidae | | | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| | Sphyraena jello | Jedd | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |

Table 10 (d) - CPUE in kg/boathour for tarad landings at Al Bateen (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|------------------------|-------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|-------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Al Sakkar</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 24.9 | 11.6 | 0.0 | 0.0 | 0.0 | 22.0 | 4.9 | 8.5 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Belonidae et al. | Haqool | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.6 | 1.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.8 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.6 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.7 |
| Chanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.8 |
| | Chanos chanos | Aifah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.8 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.3 | 10.0 | 0.0 | 0.0 | 0.0 | 9.1 | 2.0 | 3.4 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.3 | 10.0 | 0.0 | 0.0 | 0.0 | 9.1 | 2.0 | 3.4 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.4 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.4 |
| Lethrinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Lethrinus borbonicus | Yemah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Lethrinus nebulosus | Shaari | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.2 | 0.3 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.2 | 0.3 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.2 | 0.3 |
| | Portunus pelagicus | Qabqoob | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.2 | 0.3 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| | Scarus ghobban | Gain | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 8.2 | 0.7 | 1.2 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 8.2 | 0.7 | 1.2 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 0.2 | 0.4 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 0.2 | 0.4 |
| Total | | | 36.5 | 68.8 | 46.3 | 26.0 | 70.8 | 63.5 | 54.3 | 57.9 | 49.0 | 46.7 | 98.7 | 67.5 | 57.2 | 100.0 |
| (%) | | | 5.3 | 10.0 | 6.7 | 3.8 | 10.3 | 9.3 | 7.9 | 8.4 | 7.1 | 6.8 | 14.4 | 9.8 | 100.0 | |

Table 10 (e) - CPUE in kg/boathour for tarad landings at Free Port.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-------------------|-----------------------|-----------------|------------|------------|------------|--------------|------------|------------|-------------|------------|------------|------------|------------|------------|-------------|-------------|
| <i>Tarad</i> | <i>Al Defara</i> | | 0.0 | 0.0 | 7.8 | 220.5 | 0.0 | 0.0 | 97.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 27.2 | 51.0 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Belonidae et al. | Haqool | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Carangidae | | | 0.0 | 0.0 | 0.1 | 5.0 | 0.0 | 0.0 | 20.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 4.0 |
| | Carangoides bajad | Jesh Um Al Hala | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Gnathanodon speciosus | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 19.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.6 | 3.1 |
| | Scomberoides | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| | Scomberoides tol | Bassar | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.8 |
| Chanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 32.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 5.1 |
| | Chanos chanos | Aifah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 32.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 5.1 |
| Gerreidae | | | 0.0 | 0.0 | 7.4 | 170.0 | 0.0 | 0.0 | 37.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 17.9 | 33.6 |
| | Gerres longirostris | Badah | 0.0 | 0.0 | 7.4 | 170.0 | 0.0 | 0.0 | 37.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 17.9 | 33.6 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.8 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.8 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 45.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.8 | 7.0 |
| | Portunus pelagicus | Qabqoob | 0.0 | 0.0 | 0.0 | 45.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.8 | 7.0 |
| Siganidae | | | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.4 |
| | Acanthopagrus latus | Shaam | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |

Table 10 (e) - CPUE in kg/boathour for tarad landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|---------------------|--------------------------------|-----------------|------------|------------|------------|-------------|-------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Al Hadhra</i> | | 0.0 | 0.0 | 0.0 | 25.2 | 23.6 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 4.9 | 9.2 |
| Carangidae | | | 0.0 | 0.0 | 0.0 | 1.7 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.3 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 6.7 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 1.1 |
| | <i>Plectorhinchus sordidus</i> | Yanam | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 6.5 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 1.1 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 15.3 | 15.8 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.4 | 6.4 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 15.0 | 9.5 | 10.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 5.4 |
| | <i>Lutjanus fulviflamma</i> | Aqalah | 0.0 | 0.0 | 0.0 | 0.0 | 6.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 1.0 |
| | <i>Lutjanus malabaricus</i> | Hamra | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Valamugil seheli</i> | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Portunidae | | | 0.0 | 0.0 | 0.0 | 0.2 | 3.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| | <i>Portunus pelagicus</i> | Qabqoob | 0.0 | 0.0 | 0.0 | 0.2 | 3.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.6 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Scarus ghobban</i> | Gain | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sphyraenidae | | | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |

Table 10 (e) - CPUE in kg/boathour for tarad landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|----------------------------------|-----------------|------------|------------|-------------|------------|------------|-------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 16.2 | 11.8 | 22.0 | 11.1 | 9.8 | 24.0 | 16.9 | 20.4 | 23.3 | 21.7 | 18.9 | 15.5 | 17.6 | 33.1 |
| Ariidae | | | 0.2 | 0.1 | 0.0 | 0.2 | 0.0 | 2.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.4 | 0.3 | 0.7 |
| | <i>Arius thalassinus</i> | Khan | 0.2 | 0.1 | 0.0 | 0.2 | 0.0 | 2.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 1.4 | 0.3 | 0.7 |
| Carangidae | | | 1.9 | 2.6 | 10.7 | 4.7 | 3.9 | 6.0 | 5.5 | 3.1 | 6.9 | 2.0 | 7.2 | 2.5 | 4.7 | 8.9 |
| | <i>Alepes vari</i> | Gamlouh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Atule mate</i> | Durduman | 1.1 | 0.3 | 3.0 | 1.0 | 1.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 0.8 | 0.8 | 1.5 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.8 | 2.0 | 3.4 | 2.9 | 1.6 | 4.3 | 2.3 | 1.2 | 0.0 | 1.3 | 1.0 | 0.2 | 1.7 | 3.3 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 0.1 | 0.0 | 0.0 | 0.9 | 0.4 | 2.2 | 0.0 | 0.0 | 0.0 | 1.8 | 1.5 | 0.6 | 1.1 |
| | <i>Carangoides malabaricus</i> | Jesh Sal | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.7 | 0.0 | 0.0 | 3.8 | 0.0 | 2.3 | 0.0 | 0.6 | 1.2 |
| | <i>Caranx heberi</i> | Jesh Farow | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 1.0 | 0.4 | 0.0 | 0.0 | 0.2 | 0.3 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 1.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.3 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.4 | 0.3 | 0.1 | 0.0 | 0.1 | 0.1 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 2.6 | 0.0 | 0.0 | 0.2 | 1.0 | 0.9 | 0.3 | 0.1 | 0.0 | 0.0 | 0.4 | 0.8 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| Carcharhinidae | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.2 | 0.0 | 0.0 | 0.1 | 0.1 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.2 | 0.0 | 0.0 | 0.1 | 0.1 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.1 | 0.1 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.1 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 |
| Lethrinidae | | | 5.9 | 5.4 | 9.8 | 4.8 | 1.6 | 11.3 | 8.1 | 4.9 | 0.1 | 0.9 | 5.0 | 6.3 | 5.3 | 10.0 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 |
| | <i>Lethrinus lentjan</i> | Shaari Eshkheli | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Lethrinus microdon</i> | Souli | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Lethrinus nebulosus</i> | Shaari | 5.8 | 5.2 | 9.8 | 4.8 | 1.6 | 11.3 | 7.5 | 4.7 | 0.0 | 0.9 | 4.9 | 6.3 | 5.2 | 9.8 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 | 7.0 | 1.8 | 0.0 | 0.3 | 0.0 | 0.8 | 1.5 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.1 | 0.0 | 7.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 1.1 |
| | <i>Lutjanus fulviflamma</i> | Aqalah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lutjanus malabaricus</i> | Hamra | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 |
| | <i>Pinjalo pinjalo</i> | Neimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.8 | 0.0 | 0.0 | 0.0 | 0.1 | 0.3 |
| Mixed | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.1 |
| | <i>Mixed species</i> | Mekhlot | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.1 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

Table 10 (e) - CPUE in kg/boathour for tarad landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|---------------------|----------------------------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 16.2 | 11.8 | 22.0 | 11.1 | 9.8 | 24.0 | 16.9 | 20.4 | 23.3 | 21.7 | 18.9 | 15.5 | 17.6 | 33.1 |
| Scombridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 1.9 | 2.3 | 0.0 | 6.3 | 5.2 | 0.0 | 0.0 | 1.6 | 3.0 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.0 | 0.0 | 0.0 | 0.0 | 3.7 | 1.9 | 2.3 | 0.0 | 6.3 | 5.2 | 0.0 | 0.0 | 1.6 | 3.0 |
| Scyllaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Thenus orientalis</i> | Umm Ar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Sepiidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Sepia pharaonis</i> | Habbar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Serranidae | | | 8.1 | 3.7 | 1.2 | 0.9 | 0.6 | 0.6 | 0.5 | 1.5 | 0.3 | 4.0 | 4.7 | 4.1 | 2.5 | 4.7 |
| | <i>Cephalopholis hemistiktos</i> | Eshnenuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Epinephelus areolatus</i> | Semman | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Epinephelus coioides</i> | Hamour | 8.1 | 3.7 | 1.2 | 0.9 | 0.6 | 0.5 | 0.5 | 1.5 | 0.3 | 4.0 | 4.7 | 4.1 | 2.5 | 4.7 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 3.3 | 7.5 | 8.9 | 0.8 | 0.2 | 1.8 | 3.3 |
| | <i>Acanthopagrus bifasciatus</i> | Faskar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Acanthopagrus latus</i> | Shaam | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Rhabdosargus sarba</i> | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.3 | 7.5 | 8.9 | 0.8 | 0.2 | 1.7 | 3.2 |
| Sphyraenidae | | | 0.0 | 0.0 | 0.3 | 0.5 | 0.0 | 1.5 | 0.1 | 0.0 | 0.4 | 0.5 | 0.8 | 0.0 | 0.3 | 0.6 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 0.0 | 0.3 | 0.5 | 0.0 | 1.5 | 0.1 | 0.0 | 0.4 | 0.5 | 0.8 | 0.0 | 0.3 | 0.6 |

Table 10 (e) - CPUE in kg/boathour for tarad landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-------------------|--------------------------------|-------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Halaq</i> | | 28.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.4 | 4.5 |
| Scombridae | | | 28.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.4 | 4.5 |
| | <i>Scomberomorus commerson</i> | Kanaad | 28.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.4 | 4.5 |

Table 10 (e) - CPUE in kg/boathour for tarad landings at Free Port (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-------------------|----------------------------|-------------|------------|------------|------------|------------|------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <i>Tarad</i> | <i>Leikh</i> | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 13.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.1 | 2.2 |
| Gerreidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| | <i>Gerres longirostris</i> | Badah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.3 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 1.7 |
| | <i>Valamugil seheli</i> | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 10.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 1.7 |
| Total | | | 44.7 | 11.8 | 29.8 | 256.9 | 33.5 | 47.7 | 114.5 | 20.4 | 23.3 | 21.7 | 18.9 | 15.5 | 53.2 | 100.0 |
| (%) | | | 7.0 | 1.8 | 4.7 | 40.2 | 5.2 | 7.5 | 17.9 | 3.2 | 3.6 | 3.4 | 3.0 | 2.4 | 100.0 | |

Table 10 (f) - CPUE in kg/boathour for tarad landings at Al Sadar.

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|----------------------------------|-----------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 0.0 | 14.0 | 12.6 | 24.1 | 13.8 | 12.8 | 9.7 | 9.5 | 12.5 | 12.9 | 11.3 | 15.5 | 13.5 | 51.7 |
| Ariidae | | | 0.0 | 0.3 | 0.0 | 1.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.6 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.3 | 0.0 | 1.1 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.6 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Belonidae et al.</i> | Haqool | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.0 | 5.1 | 4.6 | 5.9 | 3.7 | 2.5 | 4.8 | 2.9 | 0.3 | 4.2 | 2.5 | 2.7 | 3.6 | 13.7 |
| | <i>Atule mate</i> | Durduman | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.2 | 0.7 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.0 | 0.0 | 0.2 | 5.8 | 0.5 | 2.1 | 3.4 | 2.7 | 0.1 | 1.1 | 0.0 | 0.4 | 1.5 | 5.6 |
| | <i>Carangoides fulvoguttatus</i> | Jesh Zeria | 0.0 | 3.8 | 4.4 | 0.0 | 3.2 | 0.0 | 0.0 | 0.0 | 0.0 | 2.7 | 2.3 | 2.3 | 1.7 | 6.5 |
| | <i>Gnathanodon speciosus</i> | Zuraidi | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Scomberoides tol</i> | Bassar | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 1.3 | 0.3 | 0.2 | 0.0 | 0.1 | 0.0 | 0.2 | 0.7 |
| | <i>Seriola dumerili</i> | Jib | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carcharhinidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Haemulidae | | | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 0.4 | 0.6 | 0.8 | 0.0 | 0.0 | 0.0 | 0.3 | 1.1 |
| | <i>Diagramma pictum</i> | Farsh | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.6 | 0.8 | 0.0 | 0.0 | 0.0 | 0.2 | 0.8 |
| | <i>Plectorhinchus gaterinus</i> | Hilali | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| | <i>Pomadasys argenteus</i> | Naqroor | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| Labridae | | | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| | <i>Choerodon robustus</i> | Jin | 0.0 | 0.0 | 0.0 | 0.1 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.2 |
| Lethrinidae | | | 0.0 | 3.3 | 1.1 | 8.1 | 5.6 | 5.2 | 1.7 | 3.3 | 2.9 | 2.9 | 3.8 | 7.0 | 4.1 | 15.6 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.0 | 0.3 | 0.1 | 0.0 | 0.6 | 0.0 | 0.0 | 0.1 | 0.0 | 0.3 | 0.2 | 0.0 | 0.1 | 0.6 |
| | <i>Lethrinus lentjan</i> | Shaari Eshkheli | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.1 | 0.0 | 0.2 |
| | <i>Lethrinus microdon</i> | Souli | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.1 | 0.2 |
| | <i>Lethrinus nebulosus</i> | Shaari | 0.0 | 3.0 | 1.0 | 8.1 | 4.7 | 5.2 | 1.7 | 3.2 | 2.9 | 2.1 | 3.6 | 6.9 | 3.8 | 14.7 |
| Lutjanidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 |
| | <i>Lutjanus ehrenbergii</i> | Naiser | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 |
| Mixed | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| | Mixed species | Mekhlot | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.2 |
| Rachycentridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Rachycentron canadum</i> | Sikkil | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

Table 10 (f) - CPUE in kg/boathour for tarad landings at Al Sadar (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|---------------------|---------------------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Hadaq</i> | | 0.0 | 14.0 | 12.6 | 24.1 | 13.8 | 12.8 | 9.7 | 9.5 | 12.5 | 12.9 | 11.3 | 15.5 | 13.5 | 51.7 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.3 |
| | Scarus ghobban | Gain | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.5 | 0.0 | 0.0 | 0.1 | 0.3 |
| Scombridae | | | 0.0 | 0.0 | 3.9 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 8.1 | 0.0 | 0.0 | 0.0 | 1.1 | 4.3 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Scomberomorus commerson | Kanaad | 0.0 | 0.0 | 3.9 | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 8.1 | 0.0 | 0.0 | 0.0 | 1.1 | 4.3 |
| Serranidae | | | 0.0 | 3.7 | 1.7 | 7.0 | 2.3 | 5.0 | 1.9 | 2.2 | 0.3 | 3.4 | 3.6 | 3.4 | 3.1 | 11.9 |
| | Epinephelus coioides | Hamour | 0.0 | 3.7 | 1.7 | 7.0 | 2.3 | 5.0 | 1.9 | 2.2 | 0.3 | 3.4 | 3.6 | 3.4 | 3.1 | 11.9 |
| Sparidae | | | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.5 | 0.6 | 0.2 | 0.2 | 0.6 |
| | Acanthopagrus bifasciatus | Faskar | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | Argyrops spinifer | Kofar | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Rhabdosargus sarba | Qabit | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.5 | 0.6 | 0.2 | 0.1 | 0.5 |
| Sphyraenidae | | | 0.0 | 1.5 | 1.3 | 0.0 | 0.8 | 0.0 | 0.7 | 0.0 | 0.0 | 1.4 | 0.7 | 2.2 | 0.8 | 3.0 |
| | Sphyraena jello | Jedd | 0.0 | 1.5 | 1.3 | 0.0 | 0.8 | 0.0 | 0.7 | 0.0 | 0.0 | 1.4 | 0.7 | 2.2 | 0.8 | 3.0 |

Table 10 (f) - CPUE in kg/boathour for tarad landings at Al Sadar (cont.).

| Family | Species | Arabic name | Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec | Mean | (%) |
|-----------------------|--------------------------------|-----------------|------------|-------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| <i>Tarad</i> | <i>Halaq</i> | | 0.0 | 71.6 | 53.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 14.0 | 0.0 | 0.0 | 0.0 | 12.6 | 48.3 |
| Ariidae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Arius thalassinus</i> | Khan | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Belonidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Belonidae et al.</i> | Haqool | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Carangidae | | | 0.0 | 0.2 | 11.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 1.2 | 4.6 |
| | <i>Atule mate</i> | Durduman | 0.0 | 0.0 | 4.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 1.5 |
| | <i>Carangoides bajad</i> | Jesh Um Al Hala | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 | 0.0 | 0.0 | 0.2 | 0.7 |
| | <i>Scomberoides</i> | Dhil'e | 0.0 | 0.1 | 6.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 2.3 |
| Carcharhinidae | | | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| | <i>Carcharhinidae et al.</i> | Jarjoor | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 |
| Labridae | | | 0.0 | 1.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.5 |
| | <i>Choerodon robustus</i> | Jin | 0.0 | 1.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.5 |
| Lethrinidae | | | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 0.9 | 3.3 |
| | <i>Lethrinus borbonicus</i> | Yemah | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | <i>Lethrinus nebulosus</i> | Shaari | 0.0 | 0.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 0.9 | 3.3 |
| Scombridae | | | 0.0 | 67.9 | 40.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.8 | 37.5 |
| | <i>Scomberomorus commerson</i> | Kanaad | 0.0 | 67.9 | 40.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.8 | 37.5 |
| Serranidae | | | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 0.3 | 1.1 |
| | <i>Epinephelus coioides</i> | Hamour | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 | 0.0 | 0.0 | 0.3 | 1.1 |
| Sphyraenidae | | | 0.0 | 0.8 | 2.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 1.1 |
| | <i>Sphyraena jello</i> | Jedd | 0.0 | 0.8 | 2.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 | 1.7 |
| Total | | | 0.0 | 85.6 | 66.1 | 24.1 | 13.8 | 12.8 | 9.7 | 9.5 | 26.5 | 12.9 | 11.3 | 15.5 | 26.2 | 100.0 |
| (%) | | | 0.0 | 29.7 | 23.0 | 8.4 | 4.8 | 4.5 | 3.4 | 3.3 | 9.2 | 4.5 | 3.9 | 5.4 | 100.0 | |

Table 11 - Wholesale value (AED '000) of landings by boat/gear type.

| Family | Species | Arabic name | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|-----------------------|---------------------------|-------------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|----------------|-------------|
| Ariidae | | | 16.9 | 0.0 | 0.0 | 0.0 | 45.3 | 0.0 | 0.3 | 0.0 | 62.6 | 0.1 |
| | Arius thalassinus | Khan | 16.9 | 0.0 | 0.0 | 0.0 | 45.3 | 0.0 | 0.3 | 0.0 | 62.6 | 0.1 |
| Belonidae | | | 0.4 | 0.0 | 0.0 | 0.0 | 1.1 | 0.1 | 0.1 | 0.0 | 1.7 | 0.0 |
| | Belonidae et al. | Haqool | 0.4 | 0.0 | 0.0 | 0.0 | 1.1 | 0.1 | 0.1 | 0.0 | 1.7 | 0.0 |
| Carangidae | | | 5,378.5 | 0.0 | 15.4 | 2.0 | 3,404.3 | 4.5 | 57.4 | 150.4 | 9,012.6 | 14.3 |
| | Alepes vari | Gamlouh | 9.7 | 0.0 | 0.0 | 0.0 | 1.4 | 0.0 | 0.0 | 0.0 | 11.1 | 0.0 |
| | Atule mate | Durduman | 878.3 | 0.0 | 0.0 | 0.0 | 273.8 | 0.0 | 20.3 | 0.0 | 1,172.4 | 1.9 |
| | Carangidae all | Jesh unidentified | 0.0 | 0.0 | 0.0 | 0.0 | 1.9 | 0.0 | 0.0 | 0.0 | 1.9 | 0.0 |
| | Carangoides bajad | Jesh Um Al Hala | 219.6 | 0.0 | 0.0 | 1.7 | 2,188.0 | 1.7 | 6.2 | 138.8 | 2,555.9 | 4.0 |
| | Carangoides fulvoguttatus | Jesh Zeria | 124.5 | 0.0 | 0.0 | 0.0 | 229.7 | 0.0 | 11.5 | 2.6 | 368.3 | 0.6 |
| | Carangoides malabaricus | Jesh Sal | 59.8 | 0.0 | 0.0 | 0.0 | 253.0 | 0.0 | 0.0 | 0.0 | 312.8 | 0.5 |
| | Caranx heberi | Jesh Farow | 27.0 | 0.0 | 0.0 | 0.0 | 47.3 | 0.0 | 0.0 | 0.0 | 74.3 | 0.1 |
| | Caranx ignobilis | Gufdar | 479.4 | 0.0 | 0.0 | 0.0 | 14.0 | 0.0 | 0.4 | 0.0 | 493.8 | 0.8 |
| | Gnathanodon speciosus | Zuraidi | 3,323.1 | 0.0 | 0.0 | 0.0 | 256.8 | 0.0 | 0.2 | 0.0 | 3,580.1 | 5.7 |
| | Megalaspis cordyla | Diyayuh | 1.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 |
| | Parastromateus niger | Halwayoh | 25.3 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 25.3 | 0.0 |
| | Scomberoides | Dhil'e | 20.1 | 0.0 | 0.0 | 0.2 | 36.8 | 2.9 | 17.9 | 7.4 | 85.2 | 0.1 |
| | Scomberoides tol | Bassar | 11.7 | 0.0 | 15.4 | 0.1 | 50.4 | 0.0 | 0.9 | 1.6 | 80.1 | 0.1 |
| | Seriola dumerili | Jib | 90.2 | 0.0 | 0.0 | 0.0 | 51.3 | 0.0 | 0.0 | 0.0 | 141.5 | 0.2 |
| | Seriolina nigrofasciata | Hamam | 108.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 108.4 | 0.2 |
| Carcharhinidae | | | 4.5 | 0.0 | 38.5 | 0.0 | 17.5 | 0.0 | 2.4 | 0.0 | 62.8 | 0.1 |
| | Carcharhinidae et al. | Jarjoor | 4.5 | 0.0 | 38.5 | 0.0 | 17.5 | 0.0 | 2.4 | 0.0 | 62.8 | 0.1 |
| Chanidae | | | 0.9 | 0.0 | 32.9 | 24.1 | 3.4 | 6.4 | 0.5 | 0.0 | 68.2 | 0.1 |
| | Chanos chanos | Aifah | 0.9 | 0.0 | 32.9 | 24.1 | 3.4 | 6.4 | 0.5 | 0.0 | 68.2 | 0.1 |
| Clupeidae | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Sardinella longiceps | Oumah | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Ephippidae | | | 51.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 51.5 | 0.1 |
| | Ephippidae et al. | Imad | 51.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 51.5 | 0.1 |
| Gerreidae | | | 1.6 | 0.0 | 336.4 | 58.1 | 0.7 | 64.7 | 0.0 | 93.5 | 555.0 | 0.9 |
| | Gerres longirostris | Badah | 1.6 | 0.0 | 336.4 | 58.1 | 0.7 | 64.7 | 0.0 | 93.5 | 555.0 | 0.9 |
| Haemulidae | | | 2,420.2 | 0.0 | 250.0 | 20.6 | 105.6 | 7.5 | 3.7 | 0.0 | 2,807.5 | 4.4 |
| | Diagramma pictum | Farsh | 2,273.1 | 0.0 | 0.0 | 0.0 | 86.3 | 0.0 | 3.5 | 0.0 | 2,363.0 | 3.7 |
| | Plectorhinchus gaterinus | Hilali | 72.5 | 0.0 | 0.0 | 0.0 | 3.5 | 0.0 | 0.0 | 0.0 | 76.0 | 0.1 |
| | Plectorhinchus sordidus | Yanam | 70.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.1 | 0.0 | 70.8 | 0.1 |
| | Pomadasys argenteus | Naqroor | 4.6 | 0.0 | 250.0 | 20.6 | 15.1 | 7.5 | 0.0 | 0.0 | 297.8 | 0.5 |

Table 11 - Wholesale value (AED '000) of landings by boat/gear type (cont.).

| Family | Species | Arabic name | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|--------------------------|------------------------|-----------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|-----------------|-------------|
| Labridae | | | 8.0 | 0.0 | 0.0 | 0.0 | 13.9 | 0.0 | 24.5 | 0.0 | 46.5 | 0.1 |
| | Choerodon robustus | Jin | 8.0 | 0.0 | 0.0 | 0.0 | 13.9 | 0.0 | 24.5 | 0.0 | 46.5 | 0.1 |
| Lethrinidae | | | 7,587.4 | 0.0 | 0.0 | 0.0 | 4,576.5 | 1.9 | 13.4 | 69.0 | 12,248.3 | 19.4 |
| | Lethrinus borbonicus | Yemah | 263.8 | 0.0 | 0.0 | 0.0 | 26.4 | 0.1 | 0.3 | 0.0 | 290.6 | 0.5 |
| | Lethrinus lentjan | Shaari Eshkheli | 1,357.6 | 0.0 | 0.0 | 0.0 | 25.9 | 0.0 | 0.0 | 0.0 | 1,383.6 | 2.2 |
| | Lethrinus microdon | Souli | 125.7 | 0.0 | 0.0 | 0.0 | 22.6 | 0.0 | 0.0 | 0.0 | 148.3 | 0.2 |
| | Lethrinus nebulosus | Shaari | 5,840.3 | 0.0 | 0.0 | 0.0 | 4,501.5 | 1.8 | 13.1 | 69.0 | 10,425.8 | 16.5 |
| Lutjanidae | | | 47.2 | 0.0 | 3.4 | 241.3 | 127.2 | 0.0 | 0.0 | 0.0 | 419.2 | 0.7 |
| | Lutjanus ehrenbergii | Naiser | 12.2 | 0.0 | 3.4 | 241.3 | 45.6 | 0.0 | 0.0 | 0.0 | 302.4 | 0.5 |
| | Lutjanus fulviflamma | Aqalah | 12.4 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 12.5 | 0.0 |
| | Lutjanus malabaricus | Hamra | 19.7 | 0.0 | 0.0 | 0.0 | 4.4 | 0.0 | 0.0 | 0.0 | 24.0 | 0.0 |
| | Lutjanus russelli | Umm Dhrais | 3.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.0 | 0.0 |
| | Pinjalo pinjalo | Neimi | 0.0 | 0.0 | 0.0 | 0.0 | 77.2 | 0.0 | 0.0 | 0.0 | 77.2 | 0.1 |
| Mixed | | | 17.4 | 0.0 | 0.0 | 0.6 | 10.6 | 0.0 | 0.0 | 0.0 | 28.6 | 0.0 |
| | Mixed species | Mekhlot | 17.4 | 0.0 | 0.0 | 0.6 | 10.6 | 0.0 | 0.0 | 0.0 | 28.6 | 0.0 |
| Mugilidae | | | 0.0 | 0.0 | 720.4 | 2.4 | 0.0 | 7.1 | 0.0 | 71.5 | 801.5 | 1.3 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 720.4 | 2.4 | 0.0 | 7.1 | 0.0 | 71.5 | 801.5 | 1.3 |
| Myliobatidae etc. | | | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 |
| | Myliobatidea et al. | Lugmah | 1.2 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 |
| Nemipteridae | | | 49.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 49.7 | 0.1 |
| | Scolopsis taeniatus | Ebzimi | 49.7 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 49.7 | 0.1 |
| Platycephalidae | | | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 2.8 | 3.5 | 0.0 |
| | Platycephalus indicus | Wahar | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 2.8 | 3.5 | 0.0 |
| Pomacanthidae | | | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 |
| | Pomacanthus maculosus | Anfooz | 0.8 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 |
| Portunidae | | | 312.7 | 0.0 | 582.3 | 55.8 | 0.0 | 5.5 | 0.0 | 507.1 | 1,463.5 | 2.3 |
| | Portunus pelagicus | Qabqoob | 312.7 | 0.0 | 582.3 | 55.8 | 0.0 | 5.5 | 0.0 | 507.1 | 1,463.5 | 2.3 |
| Rachycentridae | | | 264.3 | 0.0 | 0.0 | 0.0 | 8.4 | 0.0 | 0.3 | 0.0 | 272.9 | 0.4 |
| | Rachycentron canadum | Sikkil | 264.3 | 0.0 | 0.0 | 0.0 | 8.4 | 0.0 | 0.3 | 0.0 | 272.9 | 0.4 |
| Scaridae | | | 48.0 | 0.0 | 0.0 | 0.0 | 34.8 | 4.2 | 0.0 | 0.0 | 87.0 | 0.1 |
| | Scarus ghobban | Gain | 48.0 | 0.0 | 0.0 | 0.0 | 34.8 | 4.2 | 0.0 | 0.0 | 87.0 | 0.1 |
| Scombridae | | | 252.1 | 0.0 | 0.0 | 134.5 | 4,583.9 | 26.1 | 2,797.2 | 101.2 | 7,894.9 | 12.5 |
| | Euthynnus affinis | Sadah | 0.4 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Rastrelliger kanagurta | Garfah | 5.5 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 5.7 | 0.0 |
| | Scomberomorus | Kanaad | 246.2 | 0.0 | 0.0 | 134.5 | 4,583.7 | 26.1 | 2,797.2 | 101.2 | 7,888.7 | 12.5 |

Table 11 - Wholesale value (AED '000) of landings by boat/gear type (cont.).

| Family | Species | Arabic name | Lansh, Gargoor | Lansh, Hadaq | Tarad, Al Defara | Tarad, Al Hadhra | Tarad, Hadaq | Tarad, Sakhar | Tarad, Halaq | Tarad, Leikh | Total | (%) |
|-----------------------|-----------------------|----------------|-------------------|-----------------|---------------------|---------------------|-----------------|------------------|-----------------|-----------------|-----------------|--------------|
| Scyllaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 |
| | Thenus orientalis | Umm Ar Rubiyan | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 |
| Sepiidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 9.1 | 0.0 | 0.0 | 0.0 | 9.1 | 0.0 |
| | Sepia pharaonis | Habbar | 0.0 | 0.0 | 0.0 | 0.0 | 9.1 | 0.0 | 0.0 | 0.0 | 9.1 | 0.0 |
| Serranidae | | | 17,655.9 | 0.0 | 0.0 | 0.0 | 6,513.4 | 0.0 | 42.9 | 61.7 | 24,273.9 | 38.4 |
| | Cephalopholis | Eshnenuh | 0.2 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Epinephelus areolatus | Semman | 2.2 | 0.0 | 0.0 | 0.0 | 1.3 | 0.0 | 0.0 | 0.0 | 3.4 | 0.0 |
| | Epinephelus coioides | Hamour | 17,653.5 | 0.0 | 0.0 | 0.0 | 6,511.9 | 0.0 | 42.9 | 61.7 | 24,270.0 | 38.4 |
| Siganidae | | | 184.3 | 0.0 | 6.0 | 15.4 | 0.0 | 25.8 | 0.0 | 94.1 | 325.6 | 0.5 |
| | Siganus canaliculatus | Safi Arabi | 184.2 | 0.0 | 6.0 | 15.4 | 0.0 | 25.8 | 0.0 | 94.1 | 325.4 | 0.5 |
| | Siganus javus | Safi Sneifi | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 |
| Sparidae | | | 889.4 | 0.0 | 30.2 | 23.5 | 1,417.1 | 5.1 | 0.0 | 1.7 | 2,367.1 | 3.7 |
| | Acanthopagrus | Faskar | 304.9 | 0.0 | 0.0 | 0.0 | 9.0 | 0.0 | 0.0 | 0.0 | 313.9 | 0.5 |
| | Acanthopagrus latus | Shaam | 76.2 | 0.0 | 0.0 | 0.0 | 8.3 | 0.0 | 0.0 | 0.0 | 84.5 | 0.1 |
| | Argyrops spinifer | Kofar | 344.8 | 0.0 | 0.0 | 0.0 | 3.8 | 0.0 | 0.0 | 0.0 | 348.6 | 0.6 |
| | Rhabdosargus sarba | Qabit | 163.6 | 0.0 | 30.2 | 23.5 | 1,396.0 | 5.1 | 0.0 | 1.7 | 1,620.1 | 2.6 |
| Sphyraenidae | | | 11.2 | 0.0 | 0.0 | 0.0 | 235.8 | 0.0 | 21.3 | 11.5 | 279.8 | 0.4 |
| | Sphyraena jello | Jedd | 11.2 | 0.0 | 0.0 | 0.0 | 235.8 | 0.0 | 21.3 | 11.5 | 279.8 | 0.4 |
| Stromateidae | | | 11.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 11.5 | 0.0 |
| | Pampus argenteus | Zubaidi | 11.5 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 11.5 | 0.0 |
| Teraponidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Terapon jarbua | Yelly | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tetraodontidae | | | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| | Arothron stellatus | Faqel | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| Total | | | 35,216.9 | 0.0 | 2,015.6 | 578.3 | 21,115.3 | 159.4 | 2,963.9 | 1,164.4 | 63,213.8 | 100.0 |
| (%) | | | 55.7 | 0.0 | 3.2 | 0.9 | 33.4 | 0.3 | 4.7 | 1.8 | 100.0 | |

Table 12 - Wholesale value (AED '000) of landings by landing site.

| Family | Species | Arabic name | Sila | Delma | Jebel Dhanna | Marfa | Al Bateen | Free Port | Al Sadar | Ghantoot | Total | (%) |
|-----------------------|---------------------------|-------------------|--------------|----------------|--------------|--------------|--------------|----------------|--------------|------------|----------------|-------------|
| Ariidae | | | 0.1 | 1.2 | 0.0 | 22.8 | 0.0 | 33.0 | 5.4 | 0.0 | 62.6 | 0.1 |
| | Arius thalassinus | Khan | 0.1 | 1.2 | 0.0 | 22.8 | 0.0 | 33.0 | 5.4 | 0.0 | 62.6 | 0.1 |
| Belonidae | | | 0.0 | 0.1 | 0.0 | 0.5 | 0.1 | 0.4 | 0.6 | 0.0 | 1.7 | 0.0 |
| | Belonidae et al. | Haqool | 0.0 | 0.1 | 0.0 | 0.5 | 0.1 | 0.4 | 0.6 | 0.0 | 1.7 | 0.0 |
| Carangidae | | | 281.0 | 1,113.1 | 0.0 | 385.6 | 201.0 | 6,382.3 | 649.5 | 0.0 | 9,012.6 | 14.3 |
| | Alepes vari | Gamlouh | 0.0 | 1.0 | 0.0 | 0.0 | 0.0 | 10.1 | 0.0 | 0.0 | 11.1 | 0.0 |
| | Atule mate | Durduman | 0.7 | 86.2 | 0.0 | 66.6 | 1.5 | 980.6 | 36.9 | 0.0 | 1,172.4 | 1.9 |
| | Carangidae all | Jesh unidentified | 1.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.9 | 0.0 |
| | Carangoides bajad | Jesh Um Al Hala | 249.6 | 181.3 | 0.0 | 297.2 | 150.5 | 1,241.5 | 435.9 | 0.0 | 2,555.9 | 4.0 |
| | Carangoides fulvoguttatus | Jesh Zeria | 0.0 | 0.5 | 0.0 | 12.1 | 14.1 | 184.6 | 157.0 | 0.0 | 368.3 | 0.6 |
| | Carangoides malabaricus | Jesh Sal | 4.5 | 0.6 | 0.0 | 0.0 | 0.0 | 307.7 | 0.0 | 0.0 | 312.8 | 0.5 |
| | Caranx heberi | Jesh Farow | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 74.2 | 0.0 | 0.0 | 74.3 | 0.1 |
| | Caranx ignobilis | Gufdar | 8.7 | 178.8 | 0.0 | 0.0 | 0.0 | 306.2 | 0.0 | 0.0 | 493.8 | 0.8 |
| | Gnathanodon speciosus | Zuraidi | 8.2 | 624.0 | 0.0 | 0.8 | 0.0 | 2,945.7 | 1.4 | 0.0 | 3,580.1 | 5.7 |
| | Megalaspis cordyla | Diyayuh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 1.5 | 0.0 |
| | Parastromateus niger | Halwayoh | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 25.3 | 0.0 | 0.0 | 25.3 | 0.0 |
| | Scomberoides | Dhil'e | 0.0 | 23.3 | 0.0 | 1.9 | 17.8 | 29.5 | 12.8 | 0.0 | 85.2 | 0.1 |
| | Scomberoides tol | Bassar | 7.4 | 0.1 | 0.0 | 7.1 | 17.1 | 43.6 | 4.9 | 0.0 | 80.1 | 0.1 |
| | Seriola dumerili | Jib | 0.0 | 14.4 | 0.0 | 0.0 | 0.0 | 126.3 | 0.7 | 0.0 | 141.5 | 0.2 |
| | Seriolina nigrofasciata | Hamam | 0.0 | 2.9 | 0.0 | 0.0 | 0.0 | 105.5 | 0.0 | 0.0 | 108.4 | 0.2 |
| Carcharhinidae | | | 5.5 | 4.9 | 0.0 | 0.1 | 38.5 | 12.5 | 1.4 | 0.0 | 62.8 | 0.1 |
| | Carcharhinidae et al. | Jarjoor | 5.5 | 4.9 | 0.0 | 0.1 | 38.5 | 12.5 | 1.4 | 0.0 | 62.8 | 0.1 |
| Chanidae | | | 3.8 | 0.0 | 0.0 | 0.0 | 63.5 | 0.9 | 0.0 | 0.0 | 68.2 | 0.1 |
| | Chanos chanos | Aifah | 3.8 | 0.0 | 0.0 | 0.0 | 63.5 | 0.9 | 0.0 | 0.0 | 68.2 | 0.1 |
| Clupeidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 |
| | Sardinella longiceps | Oumah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 |
| Ephippidae | | | 0.0 | 13.3 | 0.0 | 0.0 | 0.0 | 38.2 | 0.0 | 0.0 | 51.5 | 0.1 |
| | Ephippidae et al. | Imad | 0.0 | 13.3 | 0.0 | 0.0 | 0.0 | 38.2 | 0.0 | 0.0 | 51.5 | 0.1 |
| Gerreidae | | | 0.7 | 0.0 | 0.0 | 0.0 | 552.7 | 1.6 | 0.0 | 0.0 | 555.0 | 0.9 |
| | Gerres longirostris | Badah | 0.7 | 0.0 | 0.0 | 0.0 | 552.7 | 1.6 | 0.0 | 0.0 | 555.0 | 0.9 |
| Haemulidae | | | 128.5 | 787.4 | 0.0 | 13.0 | 278.7 | 1,575.1 | 24.8 | 0.0 | 2,807.5 | 4.4 |
| | Diagramma pictum | Farsh | 128.0 | 787.0 | 0.0 | 10.6 | 0.0 | 1,423.8 | 13.6 | 0.0 | 2,363.0 | 3.7 |
| | Plectorhinchus gaterinus | Hilali | 0.0 | 0.1 | 0.0 | 2.4 | 0.0 | 72.4 | 1.1 | 0.0 | 76.0 | 0.1 |
| | Plectorhinchus sordidus | Yanam | 0.6 | 0.3 | 0.0 | 0.0 | 0.7 | 69.2 | 0.0 | 0.0 | 70.8 | 0.1 |
| | Pomadasys argenteus | Naqroor | 0.0 | 0.0 | 0.0 | 0.0 | 278.1 | 9.7 | 10.0 | 0.0 | 297.8 | 0.5 |

Table 12 - Wholesale value (AED '000) of landings by landing site (cont.).

| Family | Species | Arabic name | Sila | Delma | Jebel Dhanna | Marfa | Al Bateen | Free Port | Al Sadar | Ghantoot | Total | (%) |
|--------------------------|-------------------------|-----------------|--------------|----------------|--------------|----------------|----------------|----------------|----------------|------------|-----------------|-------------|
| Labridae | | | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 7.9 | 38.5 | 0.0 | 46.5 | 0.1 |
| | Choerodon robustus | Jin | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 7.9 | 38.5 | 0.0 | 46.5 | 0.1 |
| Lethrinidae | | | 97.0 | 1,874.2 | 0.0 | 1,005.0 | 78.3 | 8,343.1 | 850.7 | 0.0 | 12,248.3 | 19.4 |
| | Lethrinus borbonicus | Yemah | 0.7 | 10.2 | 0.0 | 10.4 | 0.3 | 257.0 | 12.0 | 0.0 | 290.6 | 0.5 |
| | Lethrinus lentjan | Shaari Eshkheli | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1,371.5 | 12.1 | 0.0 | 1,383.6 | 2.2 |
| | Lethrinus microdon | Souli | 1.0 | 48.3 | 0.0 | 1.9 | 0.0 | 93.5 | 3.7 | 0.0 | 148.3 | 0.2 |
| | Lethrinus nebulosus | Shaari | 95.4 | 1,815.7 | 0.0 | 992.6 | 78.0 | 6,621.1 | 823.0 | 0.0 | 10,425.8 | 16.5 |
| Lutjanidae | | | 0.0 | 1.1 | 0.0 | 0.9 | 244.7 | 171.3 | 1.1 | 0.0 | 419.2 | 0.7 |
| | Lutjanus ehrenbergii | Naiser | 0.0 | 1.1 | 0.0 | 0.9 | 244.7 | 54.7 | 1.1 | 0.0 | 302.4 | 0.5 |
| | Lutjanus fulviflamma | Aqalah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 12.5 | 0.0 | 0.0 | 12.5 | 0.0 |
| | Lutjanus malabaricus | Hamra | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 24.0 | 0.0 | 0.0 | 24.0 | 0.0 |
| | Lutjanus russelli | Umm Dhrais | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.9 | 0.0 | 0.0 | 3.0 | 0.0 |
| | Pinjalo pinjalo | Neimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 77.2 | 0.0 | 0.0 | 77.2 | 0.1 |
| Mixed | | | 0.3 | 0.0 | 0.0 | 3.1 | 0.6 | 23.7 | 0.9 | 0.0 | 28.6 | 0.0 |
| | Mixed species | Mekhlot | 0.3 | 0.0 | 0.0 | 3.1 | 0.6 | 23.7 | 0.9 | 0.0 | 28.6 | 0.0 |
| Mugilidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 801.5 | 0.0 | 0.0 | 0.0 | 801.5 | 1.3 |
| | Valamugil seheli | Beyah Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 801.5 | 0.0 | 0.0 | 0.0 | 801.5 | 1.3 |
| Myliobatidae etc. | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 1.2 | 0.0 |
| | Myliobatidea et al. | Lugmah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.2 | 0.0 | 0.0 | 1.2 | 0.0 |
| Nemipteridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 49.7 | 0.0 | 0.0 | 49.7 | 0.1 |
| | Scolopsis taeniatus | Ebzimi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 49.7 | 0.0 | 0.0 | 49.7 | 0.1 |
| Platycephalidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 3.4 | 0.1 | 0.0 | 0.0 | 3.5 | 0.0 |
| | Platycephalus indicus | Wahar | 0.0 | 0.0 | 0.0 | 0.0 | 3.4 | 0.1 | 0.0 | 0.0 | 3.5 | 0.0 |
| Pomacanthidae | | | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.8 | 0.0 |
| | Pomacanthus maculosus | Anfooz | 0.0 | 0.2 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.8 | 0.0 |
| Portunidae | | | 0.0 | 9.8 | 0.0 | 0.0 | 1,150.7 | 302.9 | 0.0 | 0.0 | 1,463.5 | 2.3 |
| | Portunus pelagicus | Qabqoob | 0.0 | 9.8 | 0.0 | 0.0 | 1,150.7 | 302.9 | 0.0 | 0.0 | 1,463.5 | 2.3 |
| Rachycentridae | | | 3.6 | 10.4 | 0.0 | 0.4 | 0.0 | 256.4 | 2.1 | 0.0 | 272.9 | 0.4 |
| | Rachycentron canadum | Sikkil | 3.6 | 10.4 | 0.0 | 0.4 | 0.0 | 256.4 | 2.1 | 0.0 | 272.9 | 0.4 |
| Scaridae | | | 0.0 | 0.0 | 0.0 | 9.3 | 4.2 | 48.0 | 25.5 | 0.0 | 87.0 | 0.1 |
| | Scarus ghobban | Gain | 0.0 | 0.0 | 0.0 | 9.3 | 4.2 | 48.0 | 25.5 | 0.0 | 87.0 | 0.1 |
| Scombridae | | | 877.2 | 998.6 | 0.0 | 868.2 | 1,271.6 | 1,509.6 | 2,369.7 | 0.0 | 7,894.9 | 12.5 |
| | Euthynnus affinis | Sadah | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Rastrelliger kanagurta | Garfah | 0.0 | 4.8 | 0.0 | 0.1 | 0.0 | 0.7 | 0.1 | 0.0 | 5.7 | 0.0 |
| | Scomberomorus commerson | Kanaad | 877.2 | 993.8 | 0.0 | 868.1 | 1,271.6 | 1,508.6 | 2,369.5 | 0.0 | 7,888.7 | 12.5 |

Table 12 - Wholesale value (AED '000) of landings by landing site (cont.).

| Family | Species | Arabic name | Sila | Delma | Jebel Dhanna | Marfa | Al Bateen | Free Port | Al Sadar | Ghantoot | Total | (%) |
|-----------------------|---------------------------|----------------|--------------|----------------|--------------|----------------|--------------|-----------------|----------------|------------|-----------------|-------------|
| Scyllaridae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 | 0.0 | 6.5 | 0.0 |
| | Thenus orientalis | Umm Ar Rubiyan | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 6.5 | 0.0 | 0.0 | 6.5 | 0.0 |
| Sepiidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.1 | 0.0 | 0.0 | 9.1 | 0.0 |
| | Sepia pharaonis | Habbar | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 9.1 | 0.0 | 0.0 | 9.1 | 0.0 |
| Serranidae | | | 537.3 | 3,588.9 | 0.0 | 1,814.3 | 95.7 | 16,935.4 | 1,302.3 | 0.0 | 24,273.9 | 38.4 |
| | Cephalopholis hemistiktos | Eshnenuh | 0.0 | 0.1 | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | 0.0 | 0.4 | 0.0 |
| | Epinephelus areolatus | Semman | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 3.4 | 0.0 | 0.0 | 3.4 | 0.0 |
| | Epinephelus coioides | Hamour | 537.3 | 3,588.8 | 0.0 | 1,814.3 | 95.7 | 16,931.6 | 1,302.3 | 0.0 | 24,270.0 | 38.4 |
| Siganidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 141.2 | 184.3 | 0.0 | 0.0 | 325.6 | 0.5 |
| | Siganus canaliculatus | Safi Arabi | 0.0 | 0.0 | 0.0 | 0.0 | 141.2 | 184.2 | 0.0 | 0.0 | 325.4 | 0.5 |
| | Siganus javus | Safi Sneifi | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 | 0.1 | 0.0 |
| Sparidae | | | 0.6 | 66.9 | 0.0 | 22.1 | 414.4 | 1,820.2 | 43.0 | 0.0 | 2,367.1 | 3.7 |
| | Acanthopagrus bifasciatus | Faskar | 0.1 | 48.8 | 0.0 | 2.5 | 0.0 | 261.5 | 0.9 | 0.0 | 313.9 | 0.5 |
| | Acanthopagrus latus | Shaam | 0.0 | 3.2 | 0.0 | 0.0 | 0.0 | 81.3 | 0.0 | 0.0 | 84.5 | 0.1 |
| | Argyrops spinifer | Kofar | 0.0 | 0.7 | 0.0 | 2.0 | 0.0 | 344.1 | 1.8 | 0.0 | 348.6 | 0.6 |
| | Rhabdosargus sarba | Qabit | 0.4 | 14.2 | 0.0 | 17.6 | 414.4 | 1,133.3 | 40.2 | 0.0 | 1,620.1 | 2.6 |
| Sphyraenidae | | | 3.5 | 8.2 | 0.0 | 97.5 | 29.4 | 53.5 | 87.7 | 0.0 | 279.8 | 0.4 |
| | Sphyraena jello | Jedd | 3.5 | 8.2 | 0.0 | 97.5 | 29.4 | 53.5 | 87.7 | 0.0 | 279.8 | 0.4 |
| Stromateidae | | | 0.0 | 4.8 | 0.0 | 0.0 | 0.0 | 6.7 | 0.0 | 0.0 | 11.5 | 0.0 |
| | Pampus argenteus | Zubaidi | 0.0 | 4.8 | 0.0 | 0.0 | 0.0 | 6.7 | 0.0 | 0.0 | 11.5 | 0.0 |
| Teraponidae | | | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | Terapon jarbua | Yelly | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tetraodontidae | | | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| | Arothron stellatus | Faqel | 0.0 | 0.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| Total | | | 1,939.0 | 8,484.3 | 0.0 | 4,242.8 | 5,370.1 | 37,774.5 | 5,403.2 | 0.0 | 63,213.8 | 100.0 |
| (%) | | | 3.1 | 13.4 | 0.0 | 6.7 | 8.5 | 59.8 | 8.5 | 0.0 | 100.0 | |

Figure 1 – Total catch landed by landing site in the Emirate of Abu Dhabi.




Figure 2 - Total catch landed by month and boat type.




Figure 3 - Total catch of principal species families landed by boat/gear type combination.




Figure 4 (a) - Breakdown of leading gear type used on a Lansh in the Emirate of Abu Dhabi.




Figure 4 (b) - Breakdown of leading gear type used on a Tarad in the Emirate of Abu Dhabi.




Figure 5 - Total number of fishing trips by site and boat/gear type combination.




Figure 6 - Total number of fishing trips by month and boat/gear type combination.




Figure 7 (a) - Mean duration of lansh/gargoor fishing trips (\pm 95% CI) by landing site.




Figure 7 (b) - Mean duration of tarad/hadaq fishing trips (\pm 95% CI) by landing site.




Figure 8 (a) - CPUE (Kg/Trip) of principal species families from lanch/gargoor fisheries.





Figure 8 (a) - CPUE (Kg/Trip) of principal species families from lansh/gargoor fisheries (cont.).





Figure 8 (a) - CPUE (Kg/Trip) of principal species families from lansh/gargoor fisheries (cont.).




Figure 9 - Wholesale value of principal species families by boat/gear type combination.






Figure 10 - Wholesale value of principal species families by landing site.



Annex 1 - Landing sites and fishing grounds in the Emirate of Abu Dhabi.



Annex 2 - Commercial species used in the catch and effort data recording system.

| Category | Species | English Name | Arabic Name | Arabic Family | Code | |
|----------------------------------|-----------------------------------|------------------------------------|-------------------------|-----------------|------------|-----|
| Sparidae | <i>Acanthopagrus berda</i> | Picnic seabream | Shaam Aswad | Sparidae | ABE | |
| | <i>Acanthopagrus bifasciatus</i> | Two bar seabream | Faskar | Sparidae | ABI | |
| | <i>Acanthopagrus latus</i> | Yellowfin seabream | Shaam | Sparidae | ALA | |
| | <i>Argyrops spinifer</i> | King soldier bream | Kofar | Sparidae | ASP | |
| | <i>Diplodus sargus kotschy</i> | One spot seabream | Emshawah | Sparidae | DSK | |
| | <i>Rhabdosargus sarba</i> | Goldlined seabream | Qabit | Sparidae | RSA | |
| | <i>Sparidentex hasta</i> | Sobaity seabream | Subaiti | Sparidae | SHA | |
| | Other Sparidae | | Faskar ba'qien | Sparidae | SPX | |
| Sphyraenidae | <i>Sphyraena barracuda</i> | Great barracuda | Jedd Large | Sphyraenidae | SBR | |
| | <i>Sphyraena jello</i> | Pickhandle barracuda | Jedd | Sphyraenidae | SJE | |
| | <i>Sphyraena obtusata</i> | Obtuse barracuda | Ghely | Sphyraenidae | SOB | |
| | Other Sphyraenidae | | Jedd ba'qien | Sphyraenidae | SHX | |
| Carangidae (Trevally) | <i>Carangoides bajad</i> | Orange spotted trevally | Jesh Um Al Hala | Carangidae | CBA | |
| | <i>Carangoides fulvoguttatus</i> | Yellow spotted trevally | Jesh Zeria | Carangidae | CFU | |
| | <i>Carangoides malabaricus</i> | Malabar trevally | Jesh Sal | Carangidae | CML | |
| | <i>Caranx heberi</i> | Blacktip trevally | Jesh Farow | Carangidae | CHE | |
| | <i>Caranx ignobilis</i> | Giant trevally | Gufdar | Carangidae | CIG | |
| | <i>Elagatis bipinnulata</i> | Rainbow runner | Ghareebah | Carangidae | EBI | |
| | <i>Gnathanodon speciosus</i> | Golden trevally | Zuraidi | Carangidae | GSP | |
| | <i>Seriola dumerili</i> | Greater amberjack | Jib | Carangidae | SDU | |
| | <i>Seriolina nigrofasciata</i> | Blackbanded trevally | Hamam | Carangidae | SNI | |
| | Carangidae (Queenfish) | <i>Parastromateus niger</i> | Black pomfret | Halwayoh | Carangidae | PNI |
| | | <i>Scomberoides commersonianus</i> | Talang queenfish | Dhil'e | Carangidae | SCO |
| | | <i>Scomberoides lysan</i> | Doublespotted queenfish | Dhil'e Mezdawaj | Carangidae | SLY |
| | | <i>Scomberoides tol</i> | Needlescaled queenfish | Bassar | Carangidae | STO |
| <i>Trachinotus blochii</i> | | Snubnose pompano | Leaf | Carangidae | TBL | |
| Carangidae (Scad) | <i>Trachinotus mookalee</i> | Indian pompano | Tala | Carangidae | TMK | |
| | <i>Alepes vari</i> | Herring scad | Gamlouh | Carangidae | AVR | |
| | <i>Atule mate</i> | Yellow tail scad | Durduman | Carangidae | AMA | |
| | <i>Decapterus russelli</i> | Indian scad | Sima | Carangidae | DRU | |
| | <i>Megalaspis cordyla</i> | Torpedo scad | Diyayuh | Carangidae | MCO | |
| | <i>Selaroides leptolepis</i> | Yellow stripe scad | Seena | Carangidae | SLE | |
| | Other Carangidae | | Jesh ba'qien | Carangidae | CAX | |
| Clupeidae | <i>Nematalosa nasus</i> | Bloch's gizzard shad | Yawafah | Clupeidae | NNA | |
| | <i>Sardinella longiceps</i> | Indian oil sardinella | Oumah | Clupeidae | SLO | |
| | Other Clupeidae | | Oumah others | Clupeidae | CLX | |
| Gerreidae | <i>Gerres longirostris</i> | Longtail siver biddy | Badah | Gerreidae | GAC | |
| | Other Gerreidae | | Badah ba'qien | Gerreidae | GEX | |
| Haemulidae | <i>Diagramma pictum</i> | Painted sweetlips | Farsh | Haemulidae | DPI | |
| | <i>Plectorhinchus gaterinus</i> | Blackspotted rubberlip | Hilali | Haemulidae | PGA | |
| | <i>Plectorhinchus pictus</i> | Trout sweetlips | Farsh Monaqat | Haemulidae | PPI | |
| | <i>Plectorhinchus sordidus</i> | Sordid sweetlips | Yanam | Haemulidae | PSO | |
| | <i>Pomadasys argenteus</i> | Silver grunt | Naqroor | Haemulidae | PAG | |
| | Other Haemulidae | | Farsh ba'qien | Haemulidae | HEX | |
| Lethrinidae | <i>Lethrinus borbonicus</i> | Snub nose emperor | Yemah | Lethrinidae | LBA | |
| | <i>Lethrinus lentjan</i> | Pink ear emperor | Shari Eshkheli | Lethrinidae | LLE | |
| | <i>Lethrinus microdon</i> | Small tooth emperor | Souli | Lethrinidae | LMI | |
| | <i>Lethrinus nebulosus</i> | Spangled emperor | Shaari | Lethrinidae | LNE | |
| | Other Lethrinidae | | Shaari ba'qien | Lethrinidae | LEX | |
| Nemipteridae | <i>Nemipterus peronii</i> | Notched threadfin bream | Andaq | Nemipteridae | NPR | |
| | <i>Scolopsis ghanam</i> | Arabian monocle bream | Ain Shamaluh | Nemipteridae | SGA | |
| | <i>Scolopsis taeniatus</i> | Black streaked monocle bream | Ebzimi | Nemipteridae | STA | |
| | Other Nemipteridae | | Ebzimi ba'qien | Nemipteridae | NEX | |

Annex 2 - Commercial species used in the catch and effort data recording system (cont.).

| | | | | | |
|--------------------------------|----------------------------------|-------------------------|--------------------|--------------------|------------|
| Lutjanidae | <i>Lutjanus ehrenbergii</i> | Ehrenberg's snapper | Naiser | Lutjanidae | LEH |
| | <i>Lutjanus fulviflamma</i> | Black spot snapper | Aqalah | Lutjanidae | LFU |
| | <i>Lutjanus lutjanus</i> | Big eye snapper | Gaizawan | Lutjanidae | LUT |
| | <i>Lutjanus malabaricus</i> | Malabar blood snapper | Hamra | Lutjanidae | LMA |
| | <i>Lutjanus russelli</i> | Russell's snapper | Umm Dhrais | Lutjanidae | LRE |
| | <i>Pinjalo pinjalo</i> | Pinjalo | Neimi | Lutjanidae | PPN |
| | Other Lutjanidae | | Naiser ba'qien | Lutjanidae | LUX |
| Mugilidae | <i>Valamugil seheli</i> | Blue spot mullet | Beyah Arabi | Mugilidae | VSE |
| | Other Mugilidae | | Beyah others | Mugilidae | MUX |
| Scombridae | <i>Auxis thazard</i> | Frigate tuna | Tabban | Scombridae | ATZ |
| | <i>Euthynnus affinis</i> | Kawakawa | Sadah | Scombridae | EUF |
| | <i>Rastrelliger kanagurta</i> | Indian mackerel | Garfah | Scombridae | RAS |
| | <i>Scomberomorus commerson</i> | Kingfish | Kanaad | Scombridae | SCM |
| | Other Scombridae | | Kanaad ba'qien | Scombridae | TUX |
| Scaridae | <i>Scarus ghobban</i> | Yellow scale parrotfish | Gain | Scaridae | SGH |
| | <i>Scarus persicus</i> | Gulf parrotfish | Gain Arabi | Scaridae | SPE |
| | Other Scaridae | | Gain ba'qien | Scaridae | SCX |
| Serranidae | <i>Cephalopholis hemistiktos</i> | Yellow fin hind | Eshnenuh | Serranidae | CHM |
| | <i>Epinephelus areolatus</i> | Areolate grouper | Semman | Serranidae | EAR |
| | <i>Epinephelus bleekeri</i> | Dusky tail grouper | Semman Lidden | Serranidae | EBE |
| | <i>Epinephelus coioides</i> | Orange spotted grouper | Hamour | Serranidae | ECO |
| | <i>Epinephelus latifasciatus</i> | Striped grouper | Hamour Mokhatat | Serranidae | ELA |
| | Other Serranidae | | Hamour ba'qien | Serranidae | SEX |
| Siganidae | <i>Siganus canaliculatus</i> | White-spotted spinefoot | Safi Arabi | Siganidae | SCA |
| | <i>Siganus javus</i> | Streaked spinefoot | Safi Sneifi | Siganidae | SJV |
| Labridae | <i>Cheilinus lunulatus</i> | Broomtail wrasse | Yerab | Labridae | CLN |
| | <i>Choerodon robustus</i> | Robust tuskfish | Jin | Labridae | CRO |
| | <i>Thalassoma lunare</i> | Moon wrasse | Misan | Labridae | TLU |
| | Other Labridae | | Jin ba'qien | Labridae | LAX |
| Other demersals | <i>Arius thalassinus</i> | Giant sea catfish | Khan | Ariidae | ATH |
| | <i>Arothron stellatus</i> | Stellate puffer | Faqel | Tetraodontidae | ASZ |
| | <i>Bothus pantherinus</i> | Leopard flounder | Samak Mousa | Bothidae | BPA |
| | <i>Euryglossa orientalis</i> | Oriental sole | Shars | Soleidae | EOI |
| | <i>Pampus argenteus</i> | Silver pomfret | Zubaidi | Stromateidae | PAR |
| | <i>Parupeneus margaritatus</i> | Pearly goatfish | Hedi | Mullidae | PMT |
| | <i>Platycephalus indicus</i> | Bar tail flathead | Wahar | Platycephalidae | PIN |
| | <i>Pomacanthus maculosus</i> | Yellow bar angelfish | Anfooz | Pomacanthidae | PML |
| | <i>Saurida undosquamis</i> | Brush tooth lizardfish | Senyouth Abullaban | Synodontidae | SUN |
| | <i>Terapon jarbua</i> | Jarbua terapon | Yelly | Teraponidae | TJA |
| | All Balistidae | Triggerfish | Qargumbah | Balistidae | BAX |
| | All Ephippidae | Batfish | Imad | Ephippidae | EPX |
| | Other pelagics | <i>Chanos chanos</i> | Milkfish | Aifah | Chanidae |
| <i>Coryphaena hippurus</i> | | Common dolphin fish | Anfalooos | Coryphaenidae | CHP |
| <i>Istiophorus platypterus</i> | | Indo-pacific sailfish | Khail Al Bahar | Istiophoridae | IPL |
| <i>Rachycentron canadum</i> | | Cobia | Sikkil | Rachycentridae | RCA |
| All Belonidae | | Needle fish | Haqool | Belonidae | BEX |
| Crustaceans | <i>Portunus pelagicus</i> | Blue swimming crab | Qabqoob | Portunidae | PPL |
| | <i>Thenus orientalis</i> | Flathead locust lobster | Umm Ar Rubiyan | Scyllaridae | TOR |
| Sharks & Rays | All Shark species | Sharks | Jarjoor | Carcharhinidae,... | SXX |
| | All Ray species | Rays | Lugmah | Myliobatidea,... | RXX |
| | All guitarfish and saw fish | Guitarfish/Sawfish | Soos | Rhinobatidae,.... | GXX |
| Mixed | Mixed basket (assorted species) | Mix | Mekhlot | | MIX |

Light turquoise background means species belongs to a principal species family